

USPTO PATENT FULL-TEXT AND IMAGE DATABASE



(8 of 11)

United States Patent
Fischer , et al.

8,293,467
October 23, 2012

****Please see images for: (Certificate of Correction) ****

Biological specimen collection and transport system and methods of use

Abstract

Disclosed are compositions for isolating populations of nucleic acids from biological, forensic, and environmental samples. Also disclosed are methods for using these compositions as one-step formulations for killing pathogens, inactivating nucleases, and releasing polynucleotides from other cellular components within the sample, and stabilizing the nucleic acids prior to further processing or assay. The disclosed compositions safely facilitate rapid sample collection, and provide extended storage and transport of the samples at ambient or elevated temperature without contamination of the sample or degradation of the nucleic acids contained therein. This process particularly facilitates the collection of specimens from remote locations, and under conditions previously considered hostile for preserving the integrity of nucleic acids released from lysed biological samples without the need of refrigeration or freezing prior to molecular analysis.

Inventors: Fischer; Gerald W. (Bethesda, MD), Daum; Luke T. (San Antonio, TX)

Assignee: Longhorn Vaccines & Diagnostics LLC (San Antonio, TX)

Family ID: 40824967

Appl. No.: 13/332,204

Filed: December 20, 2011

Prior Publication Data

Document Identifier

US 20120100529 A1

Publication Date

Apr 26, 2012

Related U.S. Patent Documents

<u>Application Number</u>	<u>Filing Date</u>	<u>Patent Number</u>	<u>Issue Date</u>
12243949	Oct 1, 2008	8084443	
60976728	Oct 1, 2007		

Current U.S. Class: 435/5; 435/6.12
Current CPC Class: C12Q 1/6806 (20130101); C12Q 1/70 (20130101); C12Q 1/6806 (20130101); C12Q 2527/125 (20130101)
Current International Class: C12Q 1/70 (20060101); C12Q 1/68 (20060101)
Field of Search: ;435/5,6.12

References Cited [\[Referenced By\]](#)

U.S. Patent Documents

4116777	September 1978	Takatsy et al.
4196265	April 1980	Koprowski et al.
4235244	November 1980	Abele et al.
4315073	February 1982	Brown et al.
4355102	October 1982	Quash
4356170	October 1982	Jennings et al.
4372945	February 1983	Likhite
4474757	October 1984	Arnon et al.
4529702	July 1985	Bryan
4554101	November 1985	Hopp
4559231	December 1985	Bjerre et al.
4578770	March 1986	Mitani
4588680	May 1986	Bucher et al.
4596792	June 1986	Vyas
4599230	July 1986	Milich et al.
4601903	July 1986	Frasch
4608251	August 1986	Mia
4634664	January 1987	Oestberg
4634666	January 1987	Engleman et al.
4668476	May 1987	Bridgham et al.
4683195	July 1987	Mullis et al.
4683202	July 1987	Mullis
4707450	November 1987	Nason
4744982	May 1988	Hunter et al.
4746490	May 1988	Saneii

4749490	June 1988	Smyth et al.
4800159	January 1989	Mullis et al.
4803998	February 1989	Kezes et al.
4816513	March 1989	Bridgham et al.
4883750	November 1989	Whiteley et al.
4954449	September 1990	Hunter et al.
4981782	January 1991	Judd et al.
4996143	February 1991	Heller et al.
5091316	February 1992	Monthony et al.
5108927	April 1992	Dorn
5136019	August 1992	Judd et al.
5145684	September 1992	Liversidge et al.
5149653	September 1992	Roser
5163441	November 1992	Monthony et al.
5182109	January 1993	Tamura et al.
5186898	February 1993	Bridgham et al.
5187060	February 1993	Cerutti et al.
5234809	August 1993	Boom
5243030	September 1993	Judd et al.
5252458	October 1993	Liav et al.
5290686	March 1994	Kendal et al.
5316910	May 1994	Rota et al.
5399363	March 1995	Liversidge et al.
5482856	January 1996	Fell, Jr. et al.
5503841	April 1996	Doyle et al.
5543158	August 1996	Gref et al.
5545555	August 1996	Racioppi et al.
5552157	September 1996	Yagi et al.
5565213	October 1996	Nakamori et al.
5565322	October 1996	Heller
5567434	October 1996	Szoka, Jr.
5571511	November 1996	Fischer
5589174	December 1996	Okuno et al.
5627071	May 1997	Triva
5631350	May 1997	Okuno et al.
5641515	June 1997	Ramtoola
5656016	August 1997	Ogden
5663055	September 1997	Turner et al.
5679356	October 1997	Bonnem et al.

5691299	November 1997	Fabry
5697899	December 1997	Hillman et al.
5702944	December 1997	Racioppi et al.
5719020	February 1998	Liav et al.
5738868	April 1998	Shinkarenko
5741516	April 1998	Webb et al.
5766841	June 1998	Liav et al.
5770219	June 1998	Chiang et al.
5779708	July 1998	Wu
5783208	July 1998	Venkateshwaran et al.
5785975	July 1998	Parikh
5795582	August 1998	Wright
5795587	August 1998	Gao et al.
5797898	August 1998	Santini, Jr. et al.
5800810	September 1998	Doyle et al.
5849489	December 1998	Heller
5891624	April 1999	Huang
5945515	August 1999	Chomczynski
5955074	September 1999	Fischer
5958379	September 1999	Regenold et al.
6015664	January 2000	Henrickson et al.
6033673	March 2000	Clements
6060068	May 2000	Doyle et al.
6136585	October 2000	Ball et al.
6162603	December 2000	Heller
6168915	January 2001	Scholl et al.
6242582	June 2001	Reece et al.
6280928	August 2001	Scholl et al.
6306404	October 2001	LaPosta et al.
6306582	October 2001	Scholl et al.
6312395	November 2001	Tripp et al.
6376172	April 2002	Scholl et al.
6406842	June 2002	Scholl et al.
6440423	August 2002	Clements et al.
6451325	September 2002	Van Nest et al.
6458577	October 2002	Huang
6495316	December 2002	Scholl et al.
6500432	December 2002	Dalemans et al.
6503745	January 2003	Chand et al.

6534065	March 2003	Makin et al.
6572866	June 2003	Torcia
6573080	June 2003	Scholl et al.
6602510	August 2003	Fikes et al.
6603908	August 2003	Dallas et al.
6603998	August 2003	King et al.
6610293	August 2003	Fischer et al.
6610474	August 2003	Huang
6627396	September 2003	Swanson et al.
6632432	October 2003	Fischer
6680308	January 2004	Hassan
6689363	February 2004	Sette et al.
6713068	March 2004	Audonnet et al.
6720409	April 2004	Okuno et al.
6734292	May 2004	Omura et al.
6759241	July 2004	Hone et al.
6780421	August 2004	Haensler et al.
6793928	September 2004	van Scharrenburg et al.
6811971	November 2004	Klepp et al.
6855321	February 2005	Rappuoli et al.
6875600	April 2005	Scholl et al.
6881835	April 2005	Bai et al.
6893814	May 2005	Swanson et al.
6939543	September 2005	Fischer et al.
6946291	September 2005	Scholl et al.
7090853	August 2006	Kapp et al.
7223409	May 2007	Nagata et al.
7279162	October 2007	Fischer
7311671	December 2007	Jung et al.
7351413	April 2008	Page et al.
7357936	April 2008	Garcon
7361352	April 2008	Birkett et al.
7438919	October 2008	Dowling et al.
7494771	February 2009	Picard et al.
7541194	June 2009	Mink et al.
7547512	June 2009	Peiris
7648681	January 2010	Meyer et al.
7718402	May 2010	Gayral et al.
7794001	September 2010	Blackwell et al.

8080645	December 2011	Fischer et al.
8084443	December 2011	Fischer et al.
8097419	January 2012	Fischer et al.
2001/0021501	September 2001	Scholl et al.
2001/0034022	October 2001	Scholl et al.
2001/0036628	November 2001	Scholl et al.
2002/0054882	May 2002	Okuno et al.
2002/0055094	May 2002	Reece et al.
2002/0081567	June 2002	Henrickson et al.
2002/0082395	June 2002	Fischer et al.
2002/0169140	November 2002	Prendergast
2003/0119209	June 2003	Kaylor et al.
2003/0203357	October 2003	Huang
2003/0215796	November 2003	Scholl et al.
2003/0219442	November 2003	Mikayama et al.
2004/0009126	January 2004	Pilkiewicz et al.
2004/0013673	January 2004	Fischer et al.
2004/0071757	April 2004	Rolf
2004/0082549	April 2004	Jomaa
2004/0086849	May 2004	Shimasaki et al.
2004/0126789	July 2004	Park et al.
2004/0142319	July 2004	Yu et al.
2004/0170965	September 2004	Scholl et al.
2004/0214272	October 2004	La Rosa et al.
2004/0223976	November 2004	Bianchi et al.
2005/0009008	January 2005	Robinson et al.
2005/0042229	February 2005	Yang et al.
2005/0090009	April 2005	Cormier et al.
2005/0169941	August 2005	Lees
2005/0170334	August 2005	Mikayama et al.
2005/0181357	August 2005	Peiris et al.
2005/0187213	August 2005	Lang et al.
2005/0227269	October 2005	Lloyd et al.
2006/0002939	January 2006	Fischer et al.
2006/0014185	January 2006	Ollikka et al.
2006/0105468	May 2006	Winkler et al.
2006/0121468	June 2006	Allnutt et al.
2006/0286557	December 2006	Basehore et al.
2007/0102946	May 2007	Blackwell et al.

2007/0172835	July 2007	McBride et al.
2007/0196388	August 2007	Dowling et al.
2007/0202497	August 2007	Renuart et al.
2007/0202511	August 2007	Chen et al.
2007/0286871	December 2007	Hickle et al.
2008/0032921	February 2008	Alexander et al.
2008/0050737	February 2008	Arieli et al.
2008/0069821	March 2008	Yang et al.
2008/0075708	March 2008	Yu et al.
2008/0107665	May 2008	Suckow et al.
2008/0107687	May 2008	Poulet
2008/0118531	May 2008	Hoffmann et al.
2008/0139789	June 2008	Fischer
2008/0145373	June 2008	Arumugham et al.
2008/0181914	July 2008	Eichhorn
2008/0260763	October 2008	Felgner et al.
2009/0081202	March 2009	Fischer et al.
2009/0098527	April 2009	Fischer et al.
2009/0233309	September 2009	Fischer et al.
2009/0312285	December 2009	Fischer et al.
2010/0009343	January 2010	Fischer et al.
2010/0055672	March 2010	Sagbini
2010/0221822	September 2010	Fischer et al.
2011/0281754	November 2011	Fischer et al.
2012/0088231	April 2012	Fischer et al.
2012/0100529	April 2012	Fischer et al.
2012/0107799	May 2012	Daum
2012/0115126	May 2012	Fischer et al.

Foreign Patent Documents

0 313 224	Apr 1989	EP
320308	Jun 1989	EP
0 621 339	Oct 1994	EP
0 675 199	Oct 1995	EP
0 726 316	Aug 1996	EP
1 081 496	Mar 2001	EP
2150281	Oct 2000	RU
WO 92/03454	Mar 1992	WO
WO 92/16619	Oct 1992	WO

WO 94/09035	Apr 1994	WO
WO 94/17106	Aug 1994	WO
WO 95/08348	Mar 1995	WO
WO 97/05248	Feb 1997	WO
WO 97/46707	Dec 1997	WO
WO 97/46712	Dec 1997	WO
WO 97/46714	Dec 1997	WO
WO01/16163	Aug 2001	WO
WO 03/053462	Mar 2003	WO
WO 03/026567	Apr 2003	WO
WO 2004/004658	Jan 2004	WO
WO 2004/043407	May 2004	WO
WO 2004/055205	Jul 2004	WO
WO2004/002451	Aug 2004	WO
WO 2004/072270	Aug 2004	WO
WO 2004/084876	Oct 2004	WO
WO 2005/075642	Aug 2005	WO
WO 2005/085274	Sep 2005	WO
WO2006/041933	Apr 2006	WO
WO 2006/138444	Dec 2006	WO
WO 2007/051036	Mar 2007	WO
WO 2007/056266	May 2007	WO
WO2007051036	May 2007	WO
WO2007056266	May 2007	WO
WO 2007/091030	Aug 2007	WO
WO 2007/133682	Nov 2007	WO
WO2008079463	Mar 2008	WO
WO2008079463	Jul 2008	WO
WO2009085355	Jul 2009	WO
WO2009085355	Sep 2009	WO

Other References

US. Appl. No. 12/426,890, filed Sep. 17, 2009, Fischer et al. cited by other .

U.S. Appl. No. 13/328,992, Fischer et al. cited by other .

U.S. Appl. No. 12/243,949, filed Oct. 1, 2008, Fischer. cited by other .

U.S. Appl. No. 12/510,968, filed Jul. 28, 2009, Fischer. cited by other .

Buck et al. BioTechniques vol. 27, pp. 528-536, Sep. 1999. cited by other .

Characterization of Novel Influenza 2005. cited by other .

Daum, et al., Abstract and Poster--"Development of a Real Time Reverse-Transcription PCR

(RRT-PCR) Assay for Detection of Influenza a H1N1 2009 from Clinical Respiratory Specimens," Pediatric Infectious Disease Conference ESPID, Nice, France, May 5-8, 2010. cited by other .

De Silva et al. Influenza A virus (A/Nonhaburi/102/2009(H1N1)) segment 4 hemagglutinin (HA) gene, partial cds. Genbank Accession No. GQ 132184.1, submitted May 9, 2009. cited by other .

Hindiyeh et al. Journal of Clinical Microbiology, vol. 43, No. 2, pp. 589-595, Feb. 2005. cited by other .

"Adamantane Resistance Among Influenza, etc.", JAMA, Feb. 22, 2006, vol. 295, No. 8, pp. 891-894. cited by other .

Kutyavin et al. 3'-Minor groove binder-DNA probes increase sequence specificity at PCR extension temperatures. Nucleic Acid Res. (2000) vol. 28, No. 2, pp. 655-661. cited by other .

Lowe et al. A computer program for selection of oligonucleotide primers for polymerase chain reactions. Nucleic Acids Research (1990) vol. 18, No. 7, pp. 1757-1761. cited by other .

Denhart, M., and Doraiswamy, V., "Master Your PCR Domain!" Promega Notes, 78: 9-12 (2001). cited by other .

Master Your PCR Domain. cited by other .

<http://www.ncbi.nlm.nih.gov/genomes/FLU/SwineFlu2009.html>. NCBI Influenza Virus Resource "CLE I. GenBank Sequence from Pandemic (H1N1) 2009 Viruses". 1237 pages. cited by other .

Pamphlet--"Prime PCR System"--**Longhorn Vaccines** & Diagnostics. cited by other .

Magari, R.T., Assessing shelf life using real-time and accelerated stability tests, BioPharm Nov. 2003. cited by other .

"Taq PCR Master Mix (2x)," USB Corp., (2007). cited by other .

"Immunofluorescence and Fluorescent-Antibody Techniques", Tortora, et al., Microbiology--An Introduction, Fourth Edition, 1992, pp. 461-463. cited by other .

World Health Organization, "CDC protocol of realtime RTPCR for influenza A (H1N1)," Apr. 28, 2009. cited by other .

Wiecek, A., "Pocket PCR: The Whole Chain Reaction in His Hand," Biotechniques.com, Oct. 26, 2010. cited by other .

Danila Valmori et al. "Use of human universally antigenic tetanus toxin T cell epitopes as carriers for human vaccination" Journal of Immunology Jul. 15, 1992. cited by other .

PCT Search Report for PCT/US2008/074521 dated Feb. 13, 2009. cited by other .

PCT Written Opinion for PCT/US2008/074521 dated May 3, 2009. cited by other .

PCT Search Report for PCT/US10/43546 dated Nov. 16, 2010. cited by other .

PCT Search Report for PCT/US10/31716 dated Jul. 28, 2010. cited by other .

PCT Written Opinion for PCT/US10/31716 dated Oct. 25, 2011. cited by other .

De Folette et al. Vaccine 2006, Jun. 12, vol. 24, No. 44-46, pp. 6597-6601. cited by other .

Galarza et al. Viral Immunity 2005, vol. 18, No. 2, pp. 365-372. cited by other .

Arend et al. Infection and Immunity, 2000, vol. 68, No. 6, pp. 3314-3321. cited by other .

Geysen, et al., "Use of Peptide Synthesis to Probe Viral Antigens for Epitopes to a Resolution of a Single Amino Acid," Proc. Natl. Acad. Sci., 81, pp. 3998-4002 (1984). cited by other .

Valmori, et al., "Use of Human Universally Antigenic Tetanus Toxin T Cell Epitopes as Carriers for Human Vaccination," J. of Immunol., 149(2), pp. 717-721 (1992). cited by other .

Tolman, et al., "Cyclic V3-Loop Related HIV-1 Conjugate **Vaccines**," Int. J. Peptide Protein Res., 41, pp. 455-466 (1993). cited by other .

Conley, et al., "Immunogenicity of Synthetic HIV-1 Gp120 V3-Loop Peptide-Conjugate Immunogens," Vaccine, 12(5), pp. 445-451 (1994). cited by other .

Schneider, et al., "Induction of CD8+T Cells Using Heterologous Prime-Boost Immunisation

Strategies," Immunol. Rev., 170, pp. 29-38 (1999). cited by other .

Tanghe, et al., "Improved Immunogenicity and Protective Efficacy of a Tuberculosis DNA Vaccine Encoding Ag85 by Protein Boosting," Infect. and Immun., 69(5), pp. 3041-3047 (2001). cited by other .

Gonzalo, et al., "A Heterologous Prime-Boost Regime Using DNA and Recombinant Vaccinia Virus Expressing the Leishmania infantum P36/LACK Antigen Protects BALB/c Mice from Cutaneous Leishmaniasis," Vaccine, 20, pp. 1226-1231 (2002). cited by other .

Meyer, et al., "Complement-Mediated Enhancement of Antibody Function for Neutralization of Pseudotype Virus Containing Hepatitis C Virus E2 Chimeric Glycoprotein," J. of Virol., 76(5) pp. 2150-2158 (2002). cited by other .

Robinson, "New Hope for an AIDS Vaccine," Nat. Rev. Immunol., 2, pp. 239-250 (Apr. 2002). cited by other .

Lu, et al., "Multiepitope Trojan Antigen Peptide *Vaccines* for the Induction of Antitumor CTL and Th Immune Responses," J. of Immunol., 172, pp. 4575-4582 (2004). cited by other .

Westerfield, et al., "Peptides Delivered by Immunostimulating Reconstituted Influenza Virosomes," J. of Peptide Sci., 11(11), pp. 707-712 (2005). cited by other .

Gerhard, et al., "Prospects for Universal Influenza Virus," Emerging Infectious Diseases, 12(4), pp. 569-574 (Apr. 2006). cited by other .

Luo, "Structural Biology: Antiviral Drugs Fit for a Purpose," Nature, 443, pp. 37-38 (Sep. 1, 2006). cited by other .

PepTcell Ltd., "Technology," <http://www.peptcell.com/technology.aspx> (2007). cited by other .

Stoloff, et al., "Synthetic Multi-Epitope Peptides Identified in Silico Induce Protective Immunity Against Multiple Influenza Serotypes," Eur. J. of Immunol., 37(9), pp. 2441-2449 (Aug. 2, 2007). cited by other .

Depla, et al., "Rational Design of a Multiepitope Vaccine Encoding T-Lymphocyte Epitopes for Treatment of Chronic Hepatitis B Virus Infections," J. of Virol., 82(1), pp. 435-450 (Jan. 2008). cited by other .

Chien et al. J. Clin. Microbiol. 1999, vol. 37, No. 5, 1393-1397. cited by other .

Ishioka et al. J. Immunol. vol. 162, pp. 3915-3925. cited by other .

Lederman et al. Molecular Immunology 1991, vol. 28, No. 11, pp. 1171-1181. cited by other .

PCT Search Report for PCT/US2007/078025 dated Oct. 28, 2008. cited by other .

PCT Written Opinion for PCT/US2007/078025 dated Mar. 17, 2009. cited by other .

PCT Search Report for PCT/US2008/078499 dated Jul. 23, 2009. cited by other .

Morre, et al., "RNA Amplification by Nucleic Acid Sequence-Based Amplification with an Internal Standard Enables Reliable Detection of Chlamydia trachomatis in Cervical Scrapings and Urine Samples," J. of Clinical Microbiol., 34(12): 3108-3114 (1996). cited by other .

Rosenstrauss, et al., "An Internal Control for Routine Diagnostic PCR: Design, Properties, and Effect on Clinical Performance," J. of Clinical Microbiol., 36(1): 191-197 (1998). cited by other .

Daum, et al., Abstract and Poster--"Development of a Real Time Reverse-Transcription PCR (RRT-PCR) Assay for Detection of Influenza A H1N1 2009 From Clinical Respiratory Specimens," Pediatric Infectious Disease Conference ESPID, Nice, France, May 5-8, 2009. cited by other .

Daum, et al., Abstract--"A Molecular Transport Medium (MTM) for Pathogen Inactivation, Ambient Transport and Preservation of RNA From Clinical Samples," ICAAC, Boston, MA, Sep. 12-15, 2010. cited by other .

U.S. Appl. No. 12/426,890, filed Apr. 20, 2009, Fischer et al. cited by other .

U.S. Appl. No. 12/510,968, filed Jul. 28, 2009, Fischer et al. cited by other .

Chomezynski, P. And Sacchi, N., "Single-Step Method of RNA Isolation by Acid Guanidinium Thiocyanate-Phenol-Chloroform Extraction," *Anal. Biochem.*, 162:156-9 (1987). cited by other .

Tortora, et al., "Tools of Biochemistry 5A--Ways to Isolate and Purify Proteins and Other Macromolecules," *Microbiology--An Introduction*, pp. 152-155, 4.sup.th Ed., The Benjamin/Cummings Publishing Company, Inc., United States (1992). cited by other .

Ramanujam, R. et al., "Room-Temperature-Stable PCR Reagents," *Cold Spring Harbor Laboratory Press, PCR Methods and Appl.*, 3:75-76 (1993). cited by other .

Wolff, C. et al, "Single-Tube Nested PCR With Room-Temperature-Stable Reagents," *Cold Spring Harbor Laboratory Press, PCR Methods and Appl.*, 4:376-79 (1995). cited by other .

Matthews, et al., "Immunofluorescence and Fluorescent Antibody Techniques," *Biochemistry*, pp. 461-463, 2.sup.nd Ed., The Benjamin/Cummings Publishing Company, Inc., United States (1996). cited by other .

Schultz, C.L. et al., "A Lysis, Storage, and Transportation Buffer for Long-Term, Room-Temperature Preservation of Human Clinical Lymphoid Tissue Samples Yielding High Molecular Weight Genomic DNA Suitable for Molecular Diagnosis," *Am. J. Clin. Pathol.*, 111(6):748-52 (1999). cited by other .

Daum, L.T. et al., "Genetic and Antigenetic Analysis of the First A/New Caledonia/20/99-Like H1N1 Influenza Isolates Reported in the Americas," *Emerg. Infect. Dis.*, 8(4):408-12 (Apr. 2002). cited by other .

De Moreau de Gerbey. A.I. et al., "Stable Hepatitis C Virus RNA Detection by RT-PCR During Four Days Storage," *BioMed Central, BMC Infectious Diseases*, 2:22 (2002). cited by other .

Daum, L.T. et al., "A Rapid. Single-Step Multiplex Reverse Transcription-PCR Assay for the Detection of Human H1N1, H3N2 and B Influenza Viruses," *J. of Clinic. Virol.*, 25(3): 345-50 (2002). cited by other .

Spackman, E. et al., "Development of a Real-Time Reverse Transcriptase PCR Assay for Type A Influenza Virus and the Avian H5 and H7 Hemagglutinin Subtypes," *J. of Clinic. Microbiol.*, 40(9): 3256-60 (2002). cited by other .

Blacksell, S.D. et al., "The Effect of Sample Degradation and RNA Stabilization on Classical Swine Fever Virus RT-PCR and ELISA methods," *J. Virol. Methods*, 118(1):33-7 (2004). cited by other .

Fouchier, R.A.M. et al., "Characterization of a Novel Influenza A Virus Hemagglutinin Subtype (H16) Obtained From Black-Headed Gulls," *J. of Virol.*, 79(5):2814-22 (Mar. 2005). cited by other .

Pheng, O.C. et al., "Temperature Related Storage Evaluation of an RT-PCR Test Kit For The Detection of Dengue Infection in Mosquitoes," (Research Note), *Tropical Biomedicine*, 22(1):73-6 (2005). cited by other .

"USB Taq PCR Master Mix in qPCR," *USB Corporation, Tech Tips*, 207 (2005). cited by other .

Bright, R.A., et al., "Adamantane Resistance Among Influenza A Viruses Isolated Early During the 2005-2006 Influenza Season in the United States," *JAMA*, 295(8):891-4 (Feb. 22, 2006). cited by other .

Das, A. et al., "Development of an Internal Positive Control for Rapid Diagnosis of Avian Influenza Virus Infections by Real-Time Reverse Transcription-PCR with Lyophilized Reagents," *J. of Clinic. Microbiol.*, 44(9):3065-73 (Sep. 2006). cited by other .

Daum. L.T., et al., "Molecular Analysis of Isolates From Influenza B Outbreaks in the U.S. and Nepal, 2005," *Arch. Of Virol.*, 151:1863-1874 (2006). cited by other .

Lin, B. et al., "Broad-Spectrum Respiratory Tract Pathogen Identification Using Resequencing

DNA Microarrays," *Genome Res.*, 16(4): 527-35 (2006). cited by other .

Mohany, J. et al., "Multiplex RT-PCR for Detecting Nineteen Respiratory Viruses," *J. of Clinic. Virol.*, 36: S9 (2006). cited by other .

Daum, L.T. et al., "Real-Time RT-PCR Assays For Type and Subtype Detection Of Influenza A and B Viruses," *Influenza & Other Resp. Viruses*, 1(4): 167-75 (2007). cited by other .

"PCR-Ready Clear Supreme.TM.," Syntezza Bioscience Ltd., available at <http://www.syntezza.com/egt/PCR-Ready.sub.--Clear.sub.--Supreme.pdf> (2006). cited by other .

Wang, Z., et al "Identifying Influenza Viruses with Resequencing Microarrays," *Emerg. Infect. Dis.* 12(4):638-46 (2006). cited by other .

"TechNotes Newsletter," *Applied Biosystems*, 14(4):1-37 (2007). cited by other .

Daum, L.T., et al., "Poster--A Rapid, Simplified Collection-to-Detection System for Typing and Subtyping Influenza Viruses Using Real-Time RT-PCR and Culture," American Society for Microbiology (ASM) Conference on Emerging Technologies of Medical Importance for the Diagnosis of Infectious Diseases and the Detection of Pathogenic Microbes, Beijing, China, (2008). cited by other .

Daum, L.T., et al., "Abstract--Quantitation of Influenza A Virus From Nasal and Lung Tissue of Cotton Rats Using Real-Time RT-PCR and Culture," 26.sup.th Annual Meeting of the European Society for Pediatric Infectious Diseases, Graz, Austria, (2008). cited by other .

Daum, L.T., et al., "Abstract--Development and Clinical Evaluation of Rapid Real-Time RT-PCR Assays for Detection of Influenza A and B Viruses," 26.sup.th Annual Meeting of the European Society for Pediatric Infectious Diseases, Graz, Austria, (2008). cited by other .

Daum, L.T., et al., "Poster--A Novel Specimen Collection Solution for Molecular Diagnostic Applications." The Pediatric Academic Societies (PAS) Annual Meeting, Honolulu, HI (2008). cited by other .

Daum, L.T., et al., "Poster--Real-Time RT-PCR Detection of Influenza A Virus in Asymptomatic Culture-Negative Cotton Rats," The Pediatric Academic Societies (PAS) Annual Meeting, Honolulu, HI (2008). cited by other .

Daum, L.T., et al., "Real-Time RT-PCR Detection of Influenza Virus Within Symptomatic and Asymptomatic Family Members," The 48.sup.th Annual IDSA/ICAAC, Washington D.C. (2008). cited by other .

European Patent Office, "PCT Notification of Transmittal of the International Search Report and The Written Opinion of the International Searching Authority or the Declaration, PCT International Search Report, PCT Written Opinion of the International Searching Authority--Application No. PCT/US2007/078025," Nov. 13, 2008, 10 pages. cited by other .

"Abstracts--27.sup.th Annual Meeting of the European Society for Paediatric Infectious Disease, Brussels, Belgium, Jun. 9-13, 2009," *The Ped. Infect. Dis. J.*, 28(6):el, e75, e202, e229 (Jun. 2009). cited by other .

European Patent Office, "PCT Notification of Transmittal of the International Search Report and The Written Opinion of the International Searching Authority or the Declaration, PCT International Search Report, PCT Written Opinion of the International Searching Authority--Application No. PCT/US2008/078499," Aug. 4, 2009, 13 pages. cited by other .

Borns, M. et al. "Most Accurate PCR Enzyme Improved With Hot Start Feature," *Biocompare*, available at <http://www.biocompare.com/technicalarticle/212/Most-Accurate-PCR-Enzyme-Improved-With-Hot-Start-Feature-from-Stratagene.html> (last visited Aug, 24. 2009). cited by other .

"KOD Hot Start DNA Polymerase," *Novagen*, available at <http://www.emdbiosciences.com/Products/ProductDisplay.asp?catno=71086> (last visited Aug. 24,

2009). cited by other .

"R.A.P.I.D.RTM. System," Idaho Technology Inc., available at <http://www.idahotech.com/RAPID/Rapid-Water.html> (last visited Aug. 24, 2009). cited by other .

"AgPath-ID.TM. One-Step RT-PCR Kit," Applied Biosystems, available at <http://www.abion.com/techlib/prot/bp.sub.--1005.pdf> (last visited Aug. 24, 2009). cited by other .

"PCR Optimization: Reaction Conditions and Components," Applied Biosystems, Part No. 4371091, Revision C, pp. 1-6 available at

<http://www3.appliedbiosystems.com/cms/groups/mcb.sub.--marketing/document-s/generaldocuments/cms.sub.--042520.pdf> (last visited Aug. 24, 2009). cited by other .

"Single Tube PCR Kit Manual," Takara Bio Inc., Cat. #RR021, V.02.09, pp. 1-6 available at <http://www.takara-bio.us/files/manuals/TAK.sub.--RR021.sub.--TM.pdf> (last visited Aug. 24, 2009). cited by other .

"Luminex Confirms Effectiveness of xTAG.RTM. Respiratory Viral Panel for Swine Flu Surveillance," Medical News Today, available at

<http://www.medicalnewstoday.com/printerfriendlynews.php?newsid=148498> (May 1, 2009). cited by other .

"Luminex Receives FDA Clearance for an Update to the xTAG.RTM. Respiratory Viral Panel Package Insert Package Insert to Include Data from Two New Publications on 2009 Influenza A/H1N1," available at [http://phx.corporate-ir.net/phoenix.zhtml?c=79403&p=irol-newsArticle&ID=1-307416&highlight=\(Jul.14,2009\)](http://phx.corporate-ir.net/phoenix.zhtml?c=79403&p=irol-newsArticle&ID=1-307416&highlight=(Jul.14,2009)). cited by other .

Luke T. Daum et al., "Detection and Molecular Characterization of Clinical Influenza A and B Viruses from Original Nasal Wash Specimens Preserved in PrimeStore.TM.," (2008). cited by other .

Luke T. Daum et al., "Portugal Meeting Poster (Introduction, Materials and Methods, Results, Discussion)," (2008). cited by other .

A. E. Krafft et al., "Evaluation of PCR Testing of Ethanol-Fixed Nasal Swab Specimens as an Augmented Surveillance Strategy for Influenza Virus and Adenovirus Identification," *Journal of Clinical Microbiology*, vol. 43(4), Apr. 2005, pp. 1768-1775. cited by other .

Linda C. Canas, "Clinical Laboratory: Selection, Collection, and Transport of Specimens for Viral Cultures," Department of the Air Force, Air Force Institute of Operational Health (AFIOH), Epidemiological Surveillance Division, SDE 01 44/5001, Virology Procedure Manual, Sep. 2005, pp. 1-8. cited by other .

"Collecting, preserving and shipping specimens for the diagnosis of avian influenza A(H5N1) virus infection: Guide for field operations," WHO/CDS/EPR/ARO/2006.1, Oct. 2006. cited by other .

Jamie A. Blow et al., "Viral nucleic acid stabilization by RNA extraction reagent," *Journal of Virological Methods*, 150 (2008), Feb. 4, 2008, pp. 41-44. cited by other .

CA Office action for PCT/US08/78499, dated Mar. 29, 2012. cited by other .

CA Office action for PCT/US2007/078025, dated Jan. 4, 2011. cited by other .

EPO Exam Report for PCT/US2007/078025, dated Dec. 30, 2011. cited by other .

EPO Exam Report for PCT/US2007/078025, dated Aug. 26, 2010. cited by other .

EPO Exam Report for PCT/US2007/078025, dated Jul. 6, 2009. cited by other .

EPO Exam Report for PCT/US2007/078025, dated May 18, 2009. cited by other .

AU Exam Report for PCT/US2007/078025, dated Nov. 19, 2010. cited by other .

IL Exam Report for PCT/US2007/078025, dated Mar. 16, 2011. cited by other .

NZ Exam Report for PCT/US2007/078025, dated Jul. 7, 2010. cited by other .

Israel Office Action of Jul. 19, 2012. cited by other .

EPO Supplementary Search Report for PCT/US10/31761, dated Jul. 13, 2012. cited by other .
CA Office Action for PCT/US2008/078499, dated Mar. 29, 2012. cited by other .
PCT Written Opinion for PCT/US2008/078499, dated Jul. 4, 2010. cited by other .
"Monolithic Silica Extraction Tips for Sample Preparation," CP-Analytica, available at <http://cp-analytica> (last visited Oct. 25, 2010). cited by other .
Barnard, et al., "Suitability of new chlamydia transport medium for transport of herpes simplex virus," J. of Clin. Microbiol., 24(5): 692-695 (1986). cited by other .
Eroglu, et al., "Successful cyropreservation of mouse oocytes by using low concentrations of trehalose and dimethylsulfoxide," Biol. of Rep. 80:70-78 (2009). cited by other .
Gelmi, et al., "Bacterial survival in different transport media," European Congress of Clinical Microbiology and Infectious Diseases (ECCMID), May 28-31, 2000 (poster). cited by other .
Higashiyama, T., "Novel functions and applications of terhalose," Pure Appl. Chem. 74(7): 1263-1269. cited by other .
H1N1 RTPCR Primer/Probe Sets, Intergrated DNA Technologies--H1N1, available at <http://www.idtdna.com/catalog/h1n1/page1.aspx>. cited by other .
Johnson, F.B., "Transport of viral specimens," Clin. Microbiol. Rev. 3(2): 120-131 (1990). cited by other .
Sponseller, et al., "Influenza A pandemic (H1N1) 2009 virus infection in domestic cat," Emerg. Infect. Dis. (e-publication) (2010). cited by other .
PCT Patentability Report for PCT/US2010/043546, dated Jan. 31, 2012. cited by other .
PCT Search Report and Patentability Report for PCT/US2008/074521, dated Mar. 2, 2010. cited by other.

Primary Examiner: Henley, III; Raymond
Attorney, Agent or Firm: Remenick PLLC

Parent Case Text

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application is a continuation of U.S. application Ser. No. 12/243,949, Oct. 1, 1998 now U.S. Pat. No. 8,084,443 and claims the benefit of U.S. Provisional Application No. 60/976,728, filed Oct. 1, 2007, the entire contents of which is specifically incorporated herein in its entirety by express reference thereto.

Claims

What is claimed is:

1. A method for denaturing proteins, inactivating nucleases and killing pathogens in one step without degrading nucleic acid of a biological sample containing proteins, nucleases, and nucleic acid, and suspected to contain pathogens, comprising: providing a mixture containing one or more chaotropes, one or more detergents, one or more reducing agents, one or more chelators, and one or more buffers, together present in an amount sufficient to denature proteins, inactivate nucleases, kill pathogens, and not degrade nucleic acid;

contacting the biological sample with the mixture which denatures proteins, inactivate nucleases, kill pathogens, and does not degrade nucleic acid of the biological sample; and detecting the presence and identity of or absence of the pathogens in the biological sample.

2. The method of claim 1, wherein the one or more chaotropes are present in an amount from about 0.5 M to about 6 M, the one or more detergents are present in an amount from about 0.1% to about 1% (wt./vol.); the one or more reducing agents are present in an amount from about 0.5 mM to about 0.3 M; the one or more chelators are present in an amount from about 0.01 mM to about 1 mM; and the one or more buffers are present in an amount from about 1 mM to about 1 M.

3. The method of claim 2, wherein the one or more chaotropes comprise guanidine thiocyanate, guanidine isocyanate, guanidine hydrochloride, or any combination thereof.

4. The method of claim 2, wherein the one or more detergents comprise sodium dodecyl sulfate, lithium dodecyl sulfate, sodium taurodeoxycholate, sodium taurocholate, sodium glycocholate, sodium deoxycholate, sodium cholate, sodium alkylbenzene sulfonate, N-lauroyl sarcosine, or any combination thereof.

5. The method of claim 2, wherein the one or more reducing agents comprise 2-mercaptoethanol, tris(2-carboxyethyl)phosphine, dithiothreitol, dimethylsulfoxide, tris(2-carboxyethyl)phosphine, or any combination thereof.

6. The method of claim 2, wherein the one or more chelators comprise ethylene glycol tetra acetic acid, hydroxyethylethylenediaminetriacetic acid, diethylene triamine penta acetic acid, N,N-bis(carboxymethyl)glycine, ethylenediaminetetraacetic, citrate anhydrous, sodium citrate, calcium citrate, ammonium citrate, ammonium bicitrate, citric acid, diammonium citrate, ferric ammonium citrate, lithium citrate, or any combination thereof.

7. The method of claim 2, wherein the one or more buffers comprise tris(hydroxymethyl)aminomethane, citrate, 2-(N-morpholino)ethanesulfonic acid, N,N-Bis(2-hydroxyethyl)-2-aminoethanesulfonic acid, 1,3-bis(tris(hydroxymethyl)methyl amino)propane, 4-(2-hydroxyethyl)-1-piperazine ethanesulfonic acid, 3-(N-morpholine)propanesulfonic acid, bicarbonate, phosphate, or any combination thereof.

8. The method of claim 2, wherein the mixture further comprises one or more short-chain alkanols present in an amount from about 1% to about 25% (vol./vol.).

9. The method of claim 8, wherein the one or more short-chain alkanols comprise methanol, ethanol, propanol, butanol, pentanol, hexanol, or any combination thereof.

10. The method of claim 2, wherein the mixture further comprises one or more surfactants or defoaming agents present in an amount from about 0.0001% to about 0.3% (wt./vol.).

11. The method of claim 10, wherein the one or more surfactants or defoaming agents comprise silicone polymer, polysorbate, antifoam A, Tween, or any combination thereof.

12. The method of claim 2, wherein the pathogens are influenza virus particles or influenza-infected cells, bacteria that causes tuberculosis or tuberculosis-infected cells.

13. The method of claim 1, wherein the mixture further comprises added naked RNA or DNA present in an amount of about 1 pg/mL to 1 .mu.g/mL.
14. The method of claim 13, further comprising detecting the presence or absence of the added naked DNA or RNA.
15. The method of claim 2, wherein the mixture is buffered to a pH of from 5 to 7.
16. The method of claim 15, wherein the mixture is buffered to a pH of from about 6.4 to 6.9.
17. The method of claim 2, wherein the biological sample contains viral particles, virally infected cells, bacteria, bacterially infected cells, fungal organisms, animal cells, or plant cells.
18. The method of claim 17, wherein contacting comprises detectably liberating nucleic acid contained within the particles, cells or organisms of the biological sample.
19. The method of claim 18, wherein nucleic acid of the biological sample is detectable by PCR analysis.
20. The method of claim 19, wherein the nucleic acid of the biological sample remains detectable by PCR analysis after the mixture containing the biological sample is stored at a temperature of 4C or more for a period of at least 14 days.
21. The method of claim 20, wherein about 5% or less of the nucleic acids of the biological sample are degraded after the mixture containing the biological sample is stored at a temperature of 12C or more for a period of at least 14 days.
22. The method of claim 19, wherein the nucleic acid of the biological sample remains detectable by PCR analysis after the mixture containing the biological sample is stored at a temperature of 4C or more for a period of at least 30 days.
23. The method of claim 22, wherein the nucleic acid of the biological sample remains detectable by PCR analysis after the mixture containing the biological sample is stored at a temperature of 12C or more for a period of at least 30 days.
24. A method for denaturing proteins, inactivating nucleases and killing pathogens in one step without degrading nucleic acid of a biological sample containing proteins, nucleases, and nucleic acid, and suspected to contain pathogens, comprising: providing a mixture containing one or more chaotropes, one or more detergents, one or more reducing agents, one or more chelators, one or more buffers, one or more short-chain alkanols, and one or more surfactants, together present in an amount sufficient to denature proteins, inactivate nucleases, kill pathogens, and not degrade nucleic acid; contacting the biological sample with the mixture which denatures proteins, inactivate nucleases, kill pathogens, and does not degrade nucleic acid of the biological sample, and detecting the presence and identity of or absence of the pathogens in the biological sample, wherein, said one or more chaotropes are selected from the group consisting of guanidine thiocyanate, guanidine isocyanate, guanidine hydrochloride, and any combination thereof; said one or more detergents are selected from the group consisting of sodium dodecyl sulfate, lithium dodecyl sulfate, sodium taurodeoxycholate, sodium taurocholate, sodium glycocholate, sodium deoxycholate, sodium cholate, sodium alkylbenzene sulfonate, N-lauroyl sarcosine, and any combination thereof; said one or more chelators are

selected from the group consisting of ethylene glycol tetra acetic acid, hydroxyethylethylenediaminetriacetic acid, diethylene triamine penta acetic acid, N,N-bis(carboxymethyl)glycine, ethylenediaminetetraacetic, citrate anhydrous, sodium citrate, calcium citrate, ammonium citrate, ammonium bicitrate, citric acid, diammonium citrate, ferric ammonium citrate, lithium citrate, and any combination thereof; and said one or more buffers are selected from the group consisting of tris(hydroxymethyl)aminomethane, citrate, 2-(N-morpholino)ethanesulfonic acid, N,N-Bis(2-hydroxyethyl)-2-aminoethanesulfonic acid, 1,3-bis(tris(hydroxymethyl)methyl amino)propane, 4-(2-hydroxyethyl)-1-piperazine ethanesulfonic acid, 3-(N-morpholino) propanesulfonic acid, bicarbonate, phosphate, and any combination thereof; and said one or more short-chain alkanols are selected from the group consisting of methanol, ethanol, propanol, butanol, pentanol, or hexanol, and any combination thereof; which are together present in an amount sufficient to denature proteins, inactivate nucleases; kill pathogens, and not substantially degrade nucleic acid of the biological sample upon contacting the mixture.

25. The method of claim 24, wherein: the one or more chaotropes are present in an amount from about 0.5 M to about 6 M; the one or more detergents are present in an amount from about 0.1% to about 1% (wt./vol.); the one or more reducing agents are present in an amount from about 0.5 mM to about 0.3 M; the one or more chelators are present in an amount from about 0.01 mM to about 1 mM; the one or more buffers are present in an amount from about 0.0001% to about 0.3% (wt./vol.); the one or more short-chain alkanols are present in an amount from about 1 to about 25% (vol./vol.); and the one or more surfactants are present in an amount from about 0.0001% to about 0.3%.

26. The method of claim 25, wherein the mixture further contains naked RNA or DNA that has a predetermined nucleic acid sequence as an internal positive control that is detectable by PCR analysis.

27. The method of claim 25, wherein the pathogens are influenza virus particles or influenza virus-infected cells, bacteria that causes tuberculosis or tuberculosis-infected cells.

28. The method of claim 25, wherein the biological sample contains whole blood, plasma, serum, sputum, urine, stool, a white blood cell, a red blood cell, buffy coat, tears, mucus, saliva, semen, vaginal fluid, lymphatic fluid, amniotic fluid, spinal or cerebrospinal fluid, a peritoneal effusion, a pleural effusion, an exudate, a punctuate, an epithelial smear, a biopsy, a bone marrow sample, fluid from a cyst or an abscess, synovial fluid, vitreous or aqueous humor, an eye wash or aspirate, a pulmonary lavage or lung aspirate, or any combination thereof.

29. A method for denaturing proteins, inactivating nucleases and killing pathogens in one step without degrading nucleic acid of a biological sample containing proteins, nucleases, and nucleic acid comprising: providing a mixture containing one or more chaotropes, one or more detergents, one or more reducing agents, one or more chelators, and one or more buffers, together present in an amount sufficient to denature proteins, inactivate nucleases, kill pathogens, and not degrade nucleic acid; providing the biological sample suspected to contain pathogens; contacting the biological sample with the mixture which liberates nucleic acid of the pathogens for PCR analysis; PCR amplifying the nucleic acid, if present; and determining the presence or absence of the pathogens in the biological sample.

30. The method of claim 29, wherein the one or more chaotropes are present in an amount from about 0.5 M to about 6 M, the one or more detergents are present in an amount from about 0.1% to about 1% (wt./vol.); the one or more reducing agents are present in an amount from about 0.5 mM to about 0.3 M; the one or more chelators are present in an amount from about 0.01 mM to about 1 mM; the one or more buffers are present in

an amount from about 100 mM to about 1M and the pH of the mixture is about 6.2 to 6.9.

31. A method for denaturing proteins, inactivating nucleases and killing pathogens in one step without degrading nucleic acid of a biological sample containing proteins, nucleases, and nucleic acid, and suspected to contain pathogens, comprising: providing a mixture containing one or more chaotropes, one or more detergents, one or more reducing agents, one or more chelators, and one or more buffers, together present in an amount sufficient to denature proteins, inactivate nucleases, kill pathogens, and not degrade nucleic acid, wherein the one or more chaotropes are present in an amount from about 0.5 M to about 6 M, the one or more detergents are present in an amount from about 0.1% to about 1% (wt./vol.); the one or more reducing agents are present in an amount from about 0.5 mM to about 0.3 M; the one or more chelators are present in an amount from about 0.01 mM to about 1 mM; and the one or more buffers are present in an amount from about 1 mM to about 1M.

32. The method of claim 31, wherein the one or more chaotropes comprise guanidine thiocyanate, guanidine isocyanate, guanidine hydrochloride, or any combination thereof; the one or more detergents comprise sodium dodecyl sulfate, lithium dodecyl sulfate, sodium taurodeoxycholate, sodium taurocholate, sodium glycocholate, sodium deoxycholate, sodium cholate, sodium alkylbenzene sulfonate, N-lauroyl sarcosine, or any combination thereof; the one or more reducing agents comprise 2-mercaptoethanol, tris(2-carboxyethyl)phosphine, dithiothreitol, dimethylsulfoxide, tris(2-carboxyethyl)phosphine, or any combination thereof; the one or more chelators comprise ethylene glycol tetra acetic acid, hydroxyethylethylenediaminetriacetic acid, diethylene triamine penta acetic acid, N,N-bis(carboxymethyl)glycine, ethylenediaminetetraacetic, citrate anhydrous, sodium citrate, calcium citrate, ammonium citrate, ammonium bicitrate, citric acid, diammonium citrate, ferric ammonium citrate, lithium citrate, or any combination thereof; the one or more buffers comprise tris(hydroxymethyl)aminomethane, citrate, 2-(N-morpholino)ethanesulfonic acid, N,N-Bis(2-hydroxyethyl)-2-aminoethanesulfonic acid, 1,3-bis(tris(hydroxymethyl)methyl amino)propane, 4-(2-hydroxyethyl)-1-piperazine ethanesulfonic acid, 3-(N-morpholino) propanesulfonic acid, bicarbonate, phosphate, or any combination thereof; and the mixture is buffered to a pH of from about 5 to 7.

33. The method of claim 31, wherein the mixture is buffered to a pH of from about 6.4 to 7.0.

34. The method of claim 31, further comprising contacting the biological sample with the mixture which denatures proteins, inactivate nucleases, kill pathogens, and does not degrade nucleic acid of the biological sample.

35. The method of claim 34, wherein contact with the composition is performed in one vessel and, after contact, the biological sample is non-infectious for transport.

36. The method of claim 35, wherein the contacted biological sample does not require refrigeration for subsequent detection of the presence or absence of pathogens.

37. The method of claim 36, wherein detection of the presence and identity of or absence of the pathogens can be determined for at least 7 days after contact with the mixture.

38. The method of claim 35, wherein the contacted biological sample is at an ambient temperature for a period of time prior to the detection of the presence or absence of pathogens.

39. The method of claim 38, wherein the period of time is at least 48 hours.

40. The method of claim 31, wherein the mixture further comprises one or more short-chain alkanols present in an amount from about 1% to about 25% (vol./vol.); one or more surfactants or defoaming agents present in an amount from about 0.0001% to about 0.3% (wt./vol.); and added naked RNA or DNA present in an amount of about 1 pg/mL to 1 .mu.g/mL.

41. The method of claim 40, wherein the one or more short-chain alkanols comprise methanol, ethanol, propanol, butanol, pentanol, hexanol, or any combination thereof; and the one or more surfactants or defoaming agents comprise silicone polymer, polysorbate, antifoam A, Tween, or any combination thereof.

42. The method of claim 40, further comprising detecting the presence or absence of the added naked DNA or RNA.

Description

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

Not Applicable.

NAMES OF THE PARTIES TO A JOINT RESEARCH AGREEMENT

Not Applicable.

FIELD OF THE INVENTION

The invention relates to aqueous compositions for collection, transport, and storage of a biological specimen containing a population of nucleic acids in a single reaction vessel, which can then be purified and/or analyzed using conventional molecular biology methods. In particular, the invention is directed to a one-step composition that a) inactivates viruses or microbes in the sample, b) lyses the biological cells or tissues to free the nucleic acids from cellular debris and extraneous biomolecules, c) protects the nucleic acids from degradation by endonuclease activity, and d) preserves the nucleic acids for subsequent isolation, detection, amplification, and/or molecular analysis. In a particularly advantageous application, all four functions may be achieved in a single composition, and in a single reaction vessel, and the resultant sample may be stored at ambient temperature for extended periods without significant degradation of the polynucleotides contained within the sample.

BACKGROUND OF THE INVENTION

Description of Related Art

In the field of molecular and diagnostic analysis, the ability to keep nucleic acids in a biological sample stable, whether the specimen is taken in a remote field location, a doctor's office or in a laboratory, often determines whether the nucleic acids can be successfully analyzed. Nucleic acids in a biological sample quickly degrade and/or denature at room temperature and must generally be stored under freezing

temperatures to remain stable; however, some degree of degradation still occurs over time. This problem is magnified when a specimen is collected at a remote field site, or a significant distance from a doctor's office or laboratory environment, and especially where there may be limited, or no, access to consistent and constant cooler/refrigerator/freezer conditions until the sample is analyzed, such as where access to power (i.e., electricity), or freezer equipment is not constant or is non-existent. The problem is yet further magnified when the desired nucleic acids for downstream analysis include ribonucleic acid (RNA), which is particularly susceptible to degradation, e.g., by endogenous or exogenous endonuclease activity. Specimen transport technology presently available in the art often uses special transport media for biological samples for transport to a laboratory, in particular, packaging that imposes short time, low temperature, and practicality limits.

In addition to concerns regarding specimen stability, often there are additional concerns regarding the reagents that are used to store and/or transport the collected samples. For example, the reagents themselves frequently require cold temperatures or other special care to maintain stability. Due to these stability issues, for example, transport of the reagents to a field site, storage at the field site before use, and transport of the biological specimens and reagents back to a testing site is a primary concern.

Another significant concern when working with biological specimens is the potential inoculation, release, or dissemination of live infectious pathogens or biological agents from the specimen into the environment. Specific protocols currently exist that are employed when handling samples that may be infectious or otherwise pose health or safety risks. If the sample is kept viable and/or biologically intact to preserve its integrity for testing, individuals involved in the collection, transfer, and testing process are potentially exposed to highly dangerous contagions. Additionally, innocent bystanders nearby a field site (or nearby during transport) can be exposed if a release of the contagion occurs. As a result, the required safety measures typically increase the expense and effort required to move such samples from one location to another.

Until recently, clinical laboratory methods for pathogen detection were labor-intensive, expensive processes that required highly knowledgeable and expert scientists with specific experience. The majority of clinical diagnostic laboratories employed the use of traditional culturing methods that typically require 3 to 7 days for a viral culture--and even longer for some other bacterial targets. Furthermore, traditional culturing requires collection, transport, and laboratory propagation and handling of potentially infectious biological agents such as Ebola, avian influenza, severe acute respiratory syndrome (SARS), etc.

The field of clinical molecular diagnostics changed drastically with the advent of polymerase chain reaction (PCR) in the mid eighties, however, and shortly thereafter with real-time PCR in the mid 90's. Real-time PCR (and RT-PCR) can deliver results in hours, and the majority of modern diagnostic laboratories are transitioning away from traditional culture, and into nucleic-acid-based detection platforms, such as real-time PCR. Recent improvements in detection chemistries, such as new and improved reporting/quenching fluor, minor groove binders (MGB), and stabilized amplification reagents have paved the way for more sensitive and specific pathogen detection assays that have been proven more timely, robust, and economical than antiquated culturing methods. Advances in other nucleic acid detection strategies (in addition to real-time PCR) such as transcription-mediated amplification, ligase chain reaction (LCR), microarrays, and pathogen gene chips, have also contributed to a transition from culture vials in the clinical laboratory.

Several commercial companies (e.g., Qiagen, Roche, and bioMerieux) have developed automated nucleic acid extraction instruments, and have attempted to automate the parts of the multi-part process from sample isolation to molecular analysis. For example, the Tigris DTS.RTM. (Gen-Probe, San Diego, Calif., USA)

automates the entire detection process, and in late 2004 was FDA approved for use with Gen-Probe's APTIMA COMBO 2.RTM. assay, an FDA-approved amplified nucleic acid test (NAT) for simultaneously detecting *Chlamydia trachomatis* and *Neisseria gonorrhoeae*.

Accordingly, there is a need in the art for a safe collection, storage and transport system that maintains the integrity of the nucleic acids of even a dangerous biological specimen, typically for further molecular analysis or diagnostic testing, without the need for freezing the collected biological specimen, the collection reagents, or the collected sample in the reagents, without posing a risk to workers or innocent bystanders, and allowing for the use of less expensive and more convenient transportation methods or complicated shipping precautions.

BRIEF SUMMARY OF THE INVENTION

The present invention encompasses new and useful compositions, as well as methods of making and employing them, that may advantageously improve conventional collection, lysis, transport and storage methods for the preparation of nucleic acids from one or more biological sources. Accordingly, the present invention advantageously can provide a collection and preservation formulation to inactivate and lyse a biological specimen containing nucleic acids, and preserve nucleic acids (RNA/DNA) within the biological specimen, preferably all in a single reaction vessel, such that the integrity of the nucleic acids is at least substantially maintained, and preferably entirely maintained, so that a portion of the nucleic acids are readily available for molecular diagnostic analysis.

An additional advantage of the present invention is that the formulation can enable the separated or released nucleic acids to remain at least substantially stable, without requiring consistent and constant cooler temperatures, such as refrigeration or freezing.

The one-step formulations disclosed herein accomplish the following main functions: inactivation or killing of pathogens within the sample; lysis of cells and separation or release of nucleic acids from the cells; inactivation of endogenous or exogenous nucleases and other cellular enzymes to prevent degradation of the nucleic acids present in the sample; and facilitation of collection and handling of the sample at ambient temperatures, stabilization of the nucleic acids during subsequent transport and storage of the sample, and preservation/maintenance of the integrity of one or more polynucleotides contained with the liberated nucleic acids.

The ability to achieve all of these desirable functions in a single-step formulation, preferably in a single reaction zone or reaction vessel, is a particularly marked advantage over that presently available. Presently existing technologies do not include a single-step composition that provides for inactivation of biological components containing nucleic acids, release of nucleic acids through lysis of cells and separation or release of nucleic acids, and maintenance of the integrity of the nucleic acids. Without being bound by theory, this is in part believed to be because the process of killing the biological organism present in a sample typically results in release and activation of enzymes that degrade proteins and nucleic acids. Enzymatic degradation leads to sample destruction, which prevents analysis. The present invention, however, stabilizes and preserves the integrity of nucleic acids present in the specimen for diagnostic testing.

The one-step formulations of the present invention allow for preferably simultaneous inactivation of biological components containing nucleic acids, lysis and separation or release of nucleic acids, stabilization, and preservation. In one embodiment, some or all of the inactivation, lysis and separation or release,

stabilization, and preservation, are sequential. In a preferred embodiment, however, a majority or preferably all of these functions occur simultaneously. In all embodiments, the one-step formulation is combined with the sample to initiate these functions. This is in contrast to previous technology in which inactivation did not necessarily occur, and lysis, stabilization, and preservation occurred in a succession of separate steps, each step typically using one or more distinct reagents and protocols that were separately added.

The sequential format of prior procedures was needed to minimize errors, avoid reagent incompatibility, and provide stepwise control of results. The present invention provides all these benefits and adds the further benefits of maintaining the integrity of the nucleic acids, rendering them ready for extraction and purification, thereby improving their ultimate yield. The one-step formulation's preferably simultaneous inactivation of biological components containing nucleic acids, lysis and release of nucleic acids from cellular debris, stabilization, and preservation of nucleic acids reduces the chance for degradation of the RNA/DNA in the sample that may occur during lysis, or after lysis and before stabilization, which contributes to improved yield of the nucleic acids that are eventually extracted. An improved yield can lead to superior test results.

In one embodiment, the invention provides a composition that includes: a) one or more chaotropes (each preferably present in the composition an amount from about 0.5 M to about 6 M); b) one or more detergents (each preferably present in the composition an amount from about 0.1% to about 1%); c) one or more chelators (each preferably present in the composition in an amount from about 0.01 mM to about 1 mM); d) one or more reducing agents (each preferably present in the composition in an amount from about 0.05 M to about 0.3 M); and e) one or more defoaming agents (each preferably present in the composition in an amount from about 0.0001% to about 0.3%).

Exemplary chaotropes include, without limitation, guanidine thiocyanate (GuSCN), guanidine hydrochloride (GuHCl), guanidine isothionate, potassium thiocyanate (KSCN), sodium iodide, sodium perchlorate, urea, or any combination thereof. Descriptions of additional exemplary chaotropes and chaotropic salts can be found in, inter alia, U.S. Pat. No. 5,234,809 (specifically incorporated herein in its entirety by express reference thereto).

Exemplary detergents include, without limitation, sodium dodecyl sulfate (SDS), lithium dodecyl sulfate (LDS), sodium taurodeoxycholate (NaTDC), sodium taurocholate (NaTC), sodium glycocholate (NaGC), sodium deoxycholate (NaDC), sodium cholate, sodium alkylbenzene sulfonate (NaABS), N-lauroyl sarcosine (NLS), salts of carboxylic acids (i.e., soaps), salts of sulfonic acids, salts of sulfuric acid, phosphoric and polyphosphoric acid esters, alkylphosphates, monoalkyl phosphate (MAP), and salts of perfluorocarboxylic acids, anionic detergents including those described in U.S. Pat. No. 5,691,299 (specifically incorporated herein in its entirety by express reference thereto), or any combination thereof.

Exemplary reducing agents include, without limitation, 2-mercaptoethanol (beta.-ME), tris(2-carboxyethyl) phosphine (TCEP), dithiothreitol (DTT), formamide, dimethylsulfoxide (DMSO), or any combination thereof. In a preferred embodiment, the reducing agent includes or is TCEP.

Exemplary chelators include, without limitation, ethylene glycol tetraacetic acid (EGTA), hydroxyethylethylenediaminetriacetic acid (HEDTA), diethylene triamine pentaacetic acid (DTPA), N,N-bis(carboxymethyl)glycine (NTA), ethylenediaminetetraacetic (EDTA), citrate anhydrous, sodium citrate, calcium citrate, ammonium citrate, ammonium bicitrate, citric acid, diammonium citrate, potassium citrate, magnesium citrate, ferric ammonium citrate, lithium citrate, or any combination thereof. In preferred embodiments, the chelator includes EDTA, a citrate, or a combination thereof. In a more preferred

embodiment, the chelator includes EDT.

The compositions of the invention can further include a defoaming agent to prevent the formation of bubbles that typically result from the presence of detergents in the formulation. Defoaming agents facilitate pipetting and handling of the disclosed compositions. Exemplary surfactants/defoaming agents include, without limitation, cocoamidopropyl hydroxysultaine, alkylaminopropionic acids, imidazoline carboxylates, betaines, sulfobetaines, sultaines, alkylphenol ethoxylates, alcohol ethoxylates, polyoxyethylenated polyoxypropylene glycols, polyoxyethylenated mercaptans, long-chain carboxylic acid esters, alkonolamides, tertiary acetylenic glycols, polyoxyethylenated silicones, N-alkylpyrrolidones, alkylpolyglycosidases, silicone polymers such as Antifoam A.RTM., or polysorbates such as Tween.RTM., or any combination thereof. In a preferred embodiment, a defoaming agent includes a silicone polymer.

Optionally, the compositions of the invention may further include one or more buffers (each preferably present in the final composition in an amount from about 1 mM to about 1 M). Exemplary buffers include, without limitation, tris(hydroxymethyl)aminomethane (Tris), citrate, 2-(N-morpholino)ethanesulfonic acid (MES), N,N-Bis(2-hydroxyethyl)-2-aminoethanesulfonic Acid (BES), 1,3-bis(tris(hydroxymethyl)methylamino)propane (Bis-Tris), 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), 3-(N-morpholino)propanesulfonic acid (MOPS), N,N-bis(2-hydroxyethyl)glycine (Bicine), N-[tris(hydroxymethyl)methyl]glycine (Tricine), N-2-acetamido-2-iminodiacetic acid (ADA), N-(2-Acetamido)-2-aminoethanesulfonic acid (ACES), piperazine-1,4-bis(2-ethanesulfonic acid) (PIPES), bicarbonate, phosphate, or any combination thereof. In a preferred embodiment, the buffer includes a citrate.

The inclusion of one or more of such optional but preferred buffers is desirable to control the pH of the formulations, since it has been found that nucleic acid extraction is optimal in a pH range of about 5 to 7. Preferably, the one or more buffers employed in the disclosed compositions are chosen to provide a significant buffering capacity in the range from a pH of about 6 to a pH of about 8, more preferably within a pH range of about 6 to about 7, and more preferably still, within a pH range of about 6.2 to about 6.8. In exemplary embodiments, the pH of PrimeStore.TM. Solutions (also referred to herein as "PSS") is preferably about 6.7.+-.0.25.

The compositions of the invention may also further optionally include one or more short-chain (preferably from 1- to 6-carbon [i.e., C.sub.1-C.sub.6] alcohols) alkanols (each preferably present in the composition in an amount from about 1% to about 25%, although higher percentages of the alcohols may be employed if desired). Exemplary short-chain alkanols include linear and branched-chain alcohols, such as, without limitation, methanol, ethanol, propanol, butanol, pentanol, hexanol, or any combination thereof.

The compositions of the invention may also further optionally include one or more additional compounds or reagents including, without limitation, betaine, bovine serum albumin, and osmolytes such as trehalose, sorbitol, and the like.

In certain embodiments, the addition of nucleic acids (e.g., RNA and/or DNA) is contemplated to be beneficial for a variety of purposes and applications of the disclosed methods: a) as a "carrier" (The addition of small amounts of supplemental RNA/DNA has been previously been shown to augment/increase the overall yield of samples/specimens, particularly original specimens that may contain low amounts of target, i.e., cells, viruses, bacteria); b) as an internal positive control for downstream molecular processes and to track or monitor the fidelity of the nucleic acid preparation from sample collection to detection; and c) for comparison to a `calibrator` for downstream quantitative analysis, e.g., qRT-PCR and the like. In such

embodiments, one or more known or "control" nucleic acids could be added to the compositions in a final concentration of from about 1 pg to about 1 .mu.g.

Preferably, the compositions of the invention provide sufficient buffering capacity to adequately stabilize the populations of polynucleotides obtained from a sample, and will, most preferably, be buffered to a pH of about 6.4 to 6.9 during formulation, and will maintain the isolated populations of polynucleotides in a similar pH range when the sample is contacted with the storage/collection formulations described herein.

Preferably, the collected samples will include one or more populations of nucleic acids that are isolated from a biological sample, specimen, or source, including, for example, RNAs and DNAs.

The compositions of the present invention will typically at least substantially inactivate, and preferably entirely inactivate, any endogenous or exogenous RNAses or DNAses present in the sample, such that the nucleic acids of the sample are substantially free of any degradation, and preferably do not degrade, or lose integrity, during the collection, lysis, storage, and transport of the sample for subsequent in vitro or in vivo analyses.

Exemplary formulations of the invention include a one-step collection solution that lyses, stabilizes, and preserves the integrity of nucleic acids prepared from a biological sample for subsequent RNA and/or DNA analysis.

The disclosed compositions were developed and optimized, inter alia: 1) to facilitate preparation of high-quality nucleic acids from clinical or environmental specimens, 2) to inactivate, kill, or otherwise neutralize potentially infectious pathogens in a biological sample to facilitate safe handling and transport of the collected specimens, and 3) to stabilize released (i.e., `naked`) DNA/RNA for prolonged periods without hydrolysis or nuclease degradation of the released nucleic acids.

The compositions described herein are ideal for clinical, field and deployment use, or for high volume sample collection/extraction. Specimens collected in one or more of the disclosed compositions are biologically inactivated, and may be safely shipped, typically even without refrigeration or dry ice.

Exemplary formulations of the storage/transport/collection compositions of the invention are described in the examples herein, and include, without limitation, a composition that includes about 4 M of a chaotrope (such as guanidine thiocyanate, guanidine hydrochloride, guanidine isocyanate, or any combination thereof), about 10 mM to 30 mM of a chelator (such as EGTA, HEDTA, DTPA, NTA, EDTA, citrate anhydrous, sodium citrate, calcium citrate, ammonium citrate, ammonium bicitrate, citric acid, diammonium citrate, ferric ammonium citrate, lithium citrate, or any combination thereof), about 0.25% of a detergent (such as SDS, LDS, NaTDC, NaTC, NaGC, NaDC, sodium cholate, NaABS, NLS, or any salt or combination thereof), about 0.1 M of a reducing agent (such as .beta.-ME, DTT, DMSO, formamide, TCEP, or any combination thereof), and about 0.1% of a surfactant/defoaming agent (such as a silicone polymer [e.g., Antifoam A.RTM.] or a polysorbate [e.g., Tween.RTM.], or any combination thereof).

Additional exemplary formulations of the specimen collection compositions of the invention include, without limitation, a composition that includes about 3 M of a chaotrope (such as guanidine thiocyanate, guanidine hydrochloride, guanidine isocyanate, or any combination thereof), about 1 mM of a reducing agent (such as .beta.-ME, TCEP, formamide, DTT, DMSO, or any combination thereof), about 1 to 10 mM of a chelator (such as EGTA, HEDTA, DTPA, NTA, EDTA, citrate anhydrous, sodium citrate, calcium citrate, ammonium

citrate, ammonium bicarbonate, citric acid, diammonium citrate, ferric ammonium citrate, lithium citrate, or any combination thereof), about 0.25% of a detergent (such as SDS, LDS, NaTDC, NaTC, NaGC, NaDC, sodium cholate, NaABS, NLS, or any salt or combination thereof), and optionally but preferably about 0.0002% of a defoaming agent (also referred to as an antifoaming agent) (such as a silicone polymer or a polysorbate, or any combination thereof) and about 100 mM of a buffer (such as Tris, MES, BES, Bis-Tris, HEPES, MOPS, bicarbonate, citrate, phosphate, or any combination thereof).

Another exemplary formulation of the disclosed polynucleotide isolation and stabilization compositions include, without limitation, a composition that includes about 1 to about 4 M of a chaotropic agent such as guanidine thiocyanate, guanidine hydrochloride, or guanidine isocyanate; about 0.5 to 100 mM of a chelating agent such as EDTA, or sodium citrate, or both; about 0.1 to about 1% of an anionic detergent such as SDS or N-lauroyl sarcosine, sodium salt; about 0.001 to about 0.0001% of a surfactant or wetting agent such as the silicone polymer, Antifoam A.RTM., e); about 10 to about 500 mM of a buffering agent such as Tris-HCl; and about 10 to about 25% of a short-chain alkanol such as ethanol.

In particular embodiments, the invention provides a composition that includes about 3 M guanidine thiocyanate; about 1 mM TCEP; about 10 mM sodium citrate; about 0.5% N-lauroyl sarcosine, sodium salt; about 0.0002% Antifoam A, about 100 mM Tris-HCl, about 0.1 mM EDTA; and about 23% ethanol.

The invention also provides a method for obtaining a population of polynucleotides from a sample suspected of containing nucleic acids. The method generally involves associating the sample with an amount of one of the disclosed compositions, under conditions effective to obtain a population of polynucleotides from the sample. The invention does not require separation of the population to "obtain" the sample, as later diagnosis may or may not need such separation.

The invention also provides a method of preparing a one-step aqueous formulation of the collection/lysis/transport/storage compositions described herein for the collection of nucleic acids such as RNA and/or DNA. In an overall sense, the method generally involves combining one or more chaotropes and nuclease-free water at a temperature of about 20.degree. C. to 90.degree. C. in a reaction zone; then combining the dissolved one or more chaotropes with one or more reducing agents, one or more chelators, and one or more detergents in the reaction zone to form an intermediate composition; optionally combining a silicone polymer with the intermediate composition in an amount sufficient to minimize foaming during further preparation of the one-step aqueous formulation; combining a sufficient amount of buffer to the intermediate composition to maintain a pH of about 6 to 6.9; optionally combining a second chelating agent to the reaction zone; then increasing the temperature of the second intermediate composition to about 60 to 95.degree. C. for about 1 to 30 minutes and lowering the temperature to ambient conditions; optionally then combining a C.sub.1-6alcohol with the contents of the reaction zone; and optionally adjusting the pH to be about 6.4 to 6.9.

In additional embodiments, the invention provides a method for preparing one-step aqueous formulations adapted to obtain a population of polynucleotides from a biological sample that is suspected of containing nucleic acids. This method generally involves at least the steps of:

a) contacting the sample with an amount of the one-step aqueous formulation effective to:

i) at least substantially kill or inactivate potentially-infectious pathogens in the sample; ii) lyse a portion of cells to release RNAs and/or DNAs from the sample; and iii) substantially inhibit or prevent the released

polynucleotides in the sample from further hydrolysis or enzymatic degradation, modification, or inactivation, so as to obtain the population of polynucleotides from the sample.

Such sample may be of any origin, including, without limitation, a clinical or veterinary sample; an environmental or ecological sample, a forensic or crime scene sample, or such like, and may contain one or more nucleic acids that are of viral, microbial, animal, or plant origin, or any combination thereof.

Preferably, the methods of the invention will include at least contacting the sample with an amount of one or more of the disclosed compositions at a temperature of from 0.degree. C. to about 40.degree. C. (more preferably at a temperature of 4.degree. C. to about 35.degree. C., and still more preferably at a temperature of 10.degree. C. to about 30.degree. C.) for a period of time of at least 24 hrs, more preferably, for a period of time of at least 48 hrs, at least 72 hrs, at least 96 hrs, or longer, without causing substantial deterioration, degradation, enzymatic cleavage, and/or nucleolytic digestion, modification, or processing of the nucleic acids contained within a sample contacted with such a composition.

In certain embodiments, the methods of the invention will include at least contacting the sample with an amount of one or more of the disclosed compositions at a temperature from about 0.degree. C. to about 40.degree. C. (more preferably at a temperature from about 4.degree. C. to about 35.degree. C., still more preferably at a temperature from about 10.degree. C. to about 30.degree. C., and more preferably still at a temperature from about 15.degree. C. to about 25.degree. C.) for a period of time of at least 7 days, more preferably, for a period of time of at least 14 days, at least 21 days, at least 28 days, or even longer without causing significant deterioration, degradation, enzymatic cleavage, and/or nucleolytic processing of the nucleic acids contained within a sample so processed. It should be understood that associating a sample with an inventive composition need only occur for a short time, but to avoid the need for immediate separation of the nucleic acids from the sample and the one-step composition of the invention all the materials may remain in contact for the time periods specified above without any substantial, or without any, degradation of the nucleic acids.

Preferably, the integrity of a population of polynucleotides released from the sample into the composition will be substantially maintained, even when the composition comprising the sample is stored at ambient temperatures, and even for prolonged periods of time, including, without limitation, storage for greater than about 10 days, greater than about 20 days, or even greater than about 30 days or more. Likewise, it is desirable that the integrity of a population of polynucleotides released from the sample into the composition will be substantially maintained, even when the composition comprising the sample is stored at subtropical and tropical temperatures--even for prolonged periods of time, including, without limitation, storage for greater than about 5 days, greater than about 15 days, or even greater than about 25 days or more.

In the practice of the present methods, it is preferable that at least one or more biological cells contained within the sample are substantially lysed to release at least a first population of polynucleotides contained within such cells into the composition. Preferably, the components of the disclosed composition are sufficient to release such a population from the remaining cellular debris (including, without limitation, lipids, proteins, polysaccharides, cellular components, and such like).

It is also desirable in the practice of the present methods that at least one or more exogenous or endogenous nucleases that may be present in, on, or about the sample itself, will be sufficiently inactivated by one or more components of the composition such that the resulting nucleic acids are not destroyed, damaged, or nucleolytically cleaved when the biological cells contained within the sample are substantially lysed to

release the population of polynucleotides from the cells. Preferably, one or more components of the disclosed composition are effective to kill, inactivate, or substantially inhibit the biological activity of a DNase or an RNase, when such a protein is present in the sample.

It is also desirable in the practice of the present methods that when one or more microbes, viruses, and/or pathogens are present in, on, or about the sample when collected, such microbes, viruses, and/or pathogens will be killed or sufficiently inactivated by one or more components of the composition to facilitate safe handling of the sample by the practitioner. Preferably, one or more components of the disclosed composition are effective to render a pathogenic sample substantially, or preferably entirely, non-pathogenic without the need for adding additional components to the composition. However, in certain applications, it may also be desirable to include one or more additional anti-microbial, anti-viral, or anti-fungal agents to the compositions to render them substantially non-pathogenic, and thus, same for handling by the practitioner.

Preferably, the composition containing the sample is at least sufficiently stable, or is entirely stable, to permit storage of the sample in the composition at ambient temperature or colder at least substantially (or entirely) from the time of collection to the time of analyzing a population of polynucleotides from the sample. As used herein, "ambient temperature" can refer to temperatures of about 18.degree. C. to 25.degree. C., or in some embodiments about 20.degree. C. to 22.degree. C.

In certain embodiments, the composition containing the sample may be stored at a temperature of about 0.degree. C. to about 40.degree. C., more preferably at a temperature of about 4.degree. C. to about 30.degree. C., more preferably, at a temperature of about 10.degree. C. to about 25.degree. C., at least substantially from the time of collection to the time that the polynucleotides obtained from the sample are further isolated, purified, or characterized using one or more conventional molecular biology methodologies.

In certain embodiments, the composition containing the sample suspected of containing nucleic acids will stabilize the nucleic acids to the extent that they either remain at least substantially non-degraded (i.e., at least substantially stable) even upon prolonged storage of the composition at ambient, refrigerator, or sub-zero temperatures. It will be desirable that this stability provides that at least about 70%, at least about 85%, more preferably at least about 90%, more preferably at least about 95%, or even more preferably, at least about 98% of the polynucleotides contained within the stored sample will not be degraded upon prolonged storage of the sample. In certain embodiments, substantially all of the polynucleotides contained within the sample will be stabilized such that the original integrity of the polynucleotides is preserved during the collection, lysis, storage, and transport of the processed sample.

In certain embodiments, the method will preferably provide a population of nucleic acids prepared from a biological sample in which less than about 15% of the polynucleotides contained in the sample will be degraded during the collection, lysis, storage, and transport of the sample after it has been stored in the composition at a temperature of from -20.degree. C. to about 40.degree. C. for a period of at least 24, 48, 72, or 96 hrs or longer after the sample was initially introduced into the composition.

In related embodiments, the method will preferably provide a population of nucleic acids prepared from a biological sample in which less than about 10% of the polynucleotides contained in the sample will be degraded during the collection, lysis, storage, and transport of the sample after it has been stored in the composition at a temperature of from -20.degree. C. to about 40.degree. C. for a period of at least 24, 48, 72, or 96 hrs or longer after the sample was initially introduced into the composition.

Likewise, in some applications of the methodology disclosed herein, use of the disclosed compositions will preferably provide a population of nucleic acids that are prepared from a biological sample, wherein less than about 5% of the polynucleotides contained in the sample will be degraded during the collection, lysis, storage, and transport of the sample after it has been stored in the composition at a temperature from -20.degree. C. to about 40.degree. C. for a period of at least 24, 48, 72, or 96 hrs or longer after the sample was initially introduced into the composition.

In some instances, the population of nucleic acids prepared by the present methods may be maintained with sufficient integrity such that no more than about 1 or 2% of the sample will be degraded even when the composition is stored at a temperature from 0.degree. C. to about 40.degree. C. for periods of several days to several weeks. In fact, the inventors have shown that samples of nucleic acids isolated using the disclosed methods remain at least substantially stable, preferably stable, in their non-degraded form for periods of several weeks to even several months or more, even when the composition containing the nucleic acids is stored at a temperature from 10.degree. C. to about 40.degree. C. In one preferred embodiment, the upper limit on the above-noted temperature ranges is about 37.degree. C. Thus, the term "stable" as used herein may refer to the various embodiments noted above regarding the integrity of the population of nucleic acids after a particular time lapse at a given temperature.

Commercial Formulations and Kits

The present invention also provides kits and sample collection systems utilizing the disclosed compositions and collection/storage/transport solutions described herein. In particular embodiments, such sample collection systems may include a collection device, such as a swab, curette, or culture loop; and a collection vessel, such as a vial test tube, or specimen cup, that contains one or more of the compositions disclosed herein. The collection vessel is preferably releasably openable, such that it can be opened to insert the one-step compositions and closed and packaged, opened to insert the sample and optionally a portion of the collection device and closed for storage and transport, or both. The collection vessel may use any suitable releasably openable mechanism, including without limitation a screw cap, snap top, press-and-turn top, or the like. Such systems may also further optionally include one or more additional reagents, storage devices, transport devices, and/or instructions for obtaining, collecting, lysing, storing, or transporting samples in such systems. In a preferred embodiment, the one-step compositions of the invention may already be disposed in the reaction zone into which the sample may be associated. In such embodiments, the invention requires only a collection device and the collection vessel.

The kit may also include one or more extraction devices to help liberate and separate the nucleic acids to obtain at least substantially pure RNA/DNA to be analyzed.

Kits may also be packaged for commercial distribution, and may further optionally include one or more collection, delivery, transportation, or storage devices for sample or specimen collection, handling, or processing. The container(s) for such kits may typically include at least one vial, test tube, flask, bottle, specimen cup, or other container, into which the composition(s) may be placed, and, preferably, suitably aliquotted for individual specimen collection, transport, and storage. The kit may also include a larger container, such as a case, that includes the containers noted above, along with other equipment, instructions, and the like. The kit may also optionally include one or more additional reagents, buffers, or compounds, and may also further optionally include instructions for use of the kit in the collection of a clinical, diagnostic, environmental, or forensic sample, as well as instructions for the storage and transport of such a sample once placed in one or more of the disclosed compositions. The kit may include, e.g., multiples of about 5 or more

of the various collection devices and collection vessels and any other components to be included, so that the kits can be used to collect multiple samples from the same source or different sources.

BRIEF DESCRIPTION OF THE DRAWINGS

For promoting an understanding of the principles of the invention, reference will now be made to the embodiments, or examples, illustrated in the drawings and specific language will be used to describe the same. It will nevertheless be understood that no limitation of the scope of the invention is thereby intended. Any alterations and further modifications in the described embodiments, and any further applications of the principles of the invention as described herein are contemplated as would normally occur to one of ordinary skill in the art to which the invention relates.

The following drawings form part of the present specification and are included to demonstrate certain aspects of the present invention. The invention may be better understood by reference to the following description taken in conjunction with the accompanying drawings, in which like reference numerals identify like elements, and in which:

FIG. 1 shows the extraction efficiency of PrimeStore.TM. (ver. 1). PrimeStore (ver. 1 [depicted here as "One-step+"]) compared to the Lysis Solution provided in the RNaqueous.RTM.-Micro Kit (Ambion, Cat#AM1931) using a standard amount of whole influenza A virus. For the comparison either the one-step formulation or the Lysis Solution provided in the kit was used for viral RNA lyses and then extracted according to manufacturer protocols. Replicate reactions were processed and analyzed by real-time RT-PCR (rRT-PCR) using an ABI 7500 sequence detection system;

FIG. 2 shows the extraction efficiency of PrimeStore.TM. Solution (ver. 1) compared to commercial kits. Homogenized cotton rat nose(*) challenged with influenza A (H3N2) or a human clinical influenza A (H1N1) samples collected during the 2006-07 season were lysed in the PrimeStore.TM. Solution or lysed using the respective lyses solution, protocol, and extraction procedure from three commercially available kits: RNaqueous-Micro (Ambion Cat#AM1931), QiaAmp Viral Mini Kit (Qiagen), and AI/NCD MaxMag (Ambion) Kit. Extraction efficiency was evaluated using the ABI 7500 with the comparative CT method. The relative CT scores and viral copies detected were optimal when PrimeStore.TM. (depicted as the "one-step formulation") was utilized in place of the respective lyses buffer for each commercial kit;

FIG. 3 shows the preservation of naked RNA in PrimeStore.TM. Solution vs. Ambion RNA Storage Solution. Single-stranded Avian H5 RNA was stored in PrimeStore.TM. solution, RNA storage solution (Ambion), or water at ambient temperature (22-24.degree. C.) for 96 hours. A total of 5 pg of RNA was extracted using the RNaqueous.RTM.-Micro Kit (Ambion, Cat#AM1931) according to manufacturer recommendations and analyzed using real-time RT-PCR on an ABI 7500 (Applied Biosystems). Values are given as cycle thresholds (CT) using the absolute quantification method;

FIG. 4 shows an example of a PrimeStore.TM. packaging format for clinical diagnostic collection. (FIG. 4A). Directions of sample collection using a clinical collection swab (Copan Diagnostics) (FIG. 4B) and a 5 mL collection tube (FIG. 4C) containing 1.5 mL of PrimeStore.TM. Solution, and a schematic demonstration of the manipulation of the system (FIG. 4D and FIG. 4E).

FIG. 5 shows an exemplary commercial PrimeStore.TM. Collection Solution. Three exemplary commercial collection solution formats: 25 mL bottle, and the 5 mL and 1.5 mL tube formats;

FIG. 6 illustrates the ability of PrimeStore.TM. Solution to rapidly kill microorganisms. Shown is a comparison of cell growth of MRSA either in culture medium (TSB), or in a solution of PS. After 10 seconds in PrimeStore.TM. Solution, no viable bacterial pathogens were detected.

FIG. 7A shows the inactivation of chicken cloacal specimens in PrimeStore.TM. Solution (Ver. 1). PrimeStore.TM. Solution inactivates microbial agents in .ltoreq.1 hr. Four original chicken cloacal samples were immersed in PrimeStore.TM. Solution (top row) or water (bottom row) and subsequently plated on blood agar plates;

FIG. 7B demonstrates that PrimeStore.TM. inhibits RNA base hydrolysis for 30 days at room temperature. RNA was incubated at room temperature (22-26.degree. C.) in PrimeStore.TM. (gel lane 1 and 3) and water (gel lane 2 and 4), and subsequently RT-PCR amplified (1500 base pairs) at Day 0 and Day 30. PrimeStore.TM. preserved collected RNA, and prevented RNA/DNA degradation at room temperature up to 30 days;

FIG. 8A depicts the real-time RT-PCR analysis of "naked" influenza A avian H5 RNA template preserved in PrimeStore.TM. Solution after incubation in RNA/DNA nucleases. H5 cRNA (2 ng) was incubated with ribonuclease A and T1, and DNaseI for 1 hour @ 37.degree. C. and extracted using the RNAaqueous.RTM.-Micro Kit (Ambion). Triplicate reactions were included for each reaction condition. Real-time RT-PCR Cycle threshold (CT) values of naked RNA preserved in PrimeStore with added nucleases (average CT: 22.88) were similar to an equal quantity of template cRNA control (average CT: 23.70). Template cRNA reactions subjected to nuclease digestion without PrimeStore.TM. were almost completely degraded (average CT 39.58);

FIG. 8B depicts the real-time RT-PCR analysis and gel electrophoresis of "naked" influenza A avian H5 RNA template preserved in PrimeStore.TM. Solution after incubation in RNA/DNA nucleases @ 37.degree. C. for 7 days. Two nanograms of H5 cRNA was incubated with RNase A and T1, and DNase I, then extracted using the RNAaqueous.RTM.-Micro Kit (Ambion) after 7 days. Duplicate reactions were included for each reaction. Real-time RT-PCR Cycle Threshold (CT) values of naked RNA preserved in PrimeStore.TM. with added nucleases (average CT: 33.51) were detected after 7 days. Template cRNA reactions subjected to nuclease digestion without PrimeStore were completely degraded and similar to NTC reactions.

FIG. 8C demonstrates that PrimeStore.TM. is impervious to nuclease digestion. Gel electrophoresis of post-amplified product. Lane 3 is the PCR product from template RNA+PrimeStore.TM. at 37.degree. C. for 7 days, and Lane 5 amplification of positive control RNA. Lane 5 (no amplification) is RNA without PrimeStore.TM. Lane 2 and 6 are 100 bp ladder, and NTC reactions, respectively.

FIG. 9 illustrates that PrimeStore.TM. preservation is superior to other solutions. PrimeStore.TM. (Ver. 2 and Ver. 2.2) Preservation of RNA from influenza A virus compared to Qiagen AVL buffer, ethanol, and Viral Transport Media (VTM) at ambient temperature (22-25.degree. C.) for 30 days.

FIG. 10 shows the extraction efficiency of Influenza A virus preserved in PrimeStore.TM. (Ver. 2.2) for 30 days at various temperatures. Environmental (21-37.degree. C.); Freeze-thawed (-25.degree. C.; 32.times.); ambient temperature (22-26.degree. C.); and Lane 5: refrigerated (4.degree. C.).

FIG. 11 is a graph of critical threshold vs. molar concentration using whole influenza A virus with TCEP as

the reducing agent;

FIG. 12 is a graph of the critical threshold vs. molar concentration using H5N1 Avian influenza ssRNA with TCEP as the reducing agent;

FIG. 13A and FIG. 13B show the comparison between TCEP and .beta.-ME as reducing agent components of the PrimeStore.TM. Solution compositions, using a water only control. In FIG. 13A, H5 avian influenza RNA was employed, while in FIG. 13B, whole virus were used.

FIG. 14A shows the results of a study employing PrimeStore.TM. solution in preserving nucleic acids from blood. PrimeStore Extraction Efficiency of whole blood spiked with RNA compared to the lysis solution in the QIAamp.RTM. DNA Blood Mini Kit. 0.1 pg and 1 pg of influenza A RNA were spiked and extracted using PrimeStore.TM. or AL Lysis buffer. At both RNA concentrations, PrimeStore.TM. produced superior results as evident by real-time RT-PCR CT scores;

FIG. 14B tabulates data from the study shown in FIG. 14A involving the extraction of "naked" H5 avian influenza ssRNA from blood tubes. PrimeStore.TM. Extraction Efficiency of whole blood spiked with RNA compared to the lysis solution in the QIAamp.RTM. DNA Blood Mini Kit using different blood anticoagulants. PrimeStore.TM. was superior compared to the AL Lysis Buffer from Qiagen using Blood spiked with RNA in common anticoagulant blood-collection tubes; and

FIG. 14C tabulates data from the study shown in FIG. 14A involving the comparison of in PrimeStore.TM. Solution vs. a commercial extraction kit (Qiagen). Shown is the Extraction Efficiency of PrimeStore.TM. for whole blood spiked with RNA compared to the lysis solution in the QIAamp.RTM. DNA Blood Mini Kit. 0.1 pg and 1 pg of Influenza A viral RNA were spiked and extracted using PrimeStore or AVL Lysis buffer. At both RNA concentrations, PrimeStore.TM. produced superior results as evident by real-time RT-PCR CT scores.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

Illustrative embodiments of the invention are described below. In the interest of clarity, not all features of an actual implementation are described in this specification. It will of course be appreciated that in the development of any such actual embodiment, numerous implementation-specific decisions must be made to achieve the developers' specific goals, such as compliance with system-related and business-related constraints, which will vary from one implementation to another. Moreover, it will be appreciated that such a development effort might be complex and time-consuming, but would be a routine undertaking for those of ordinary skill in the art having the benefit of this disclosure.

The extended stabilization, collection, transport, and preservation imparted by the disclosed formulations are particularly advantageous when a sample or specimen is located in a geographical region that is remote from a testing facility. Remote locations, also referred to as field sites, encompass a variety of environments where diagnostic testing is typically not performed. These sites include doctors' offices, triage centers, airports, border crossings, outbreak areas, and a variety of outdoor locations. The disclosed compositions and methods for their use offer particular advantages in locations where there is no access to electricity and/or refrigeration, or where access is inconsistent. Because of the extended stability at room temperature, a sample can be taken from any remote location, for example without limitation at a malarial outbreak site in Africa, and the sample can be shipped to the United States or Europe for diagnostic analysis in a laboratory. Because

the disclosed collection formulations are stable at room temperature or below, and preferably even at tropical or subtropical temperatures for a time, they can routinely be taken into the field without worry that the component reagents (such as RNA controls) themselves will degrade until a sample can be analyzed, typically at a remote location from the collection.

The compositions of the invention may be any suitable aqueous formulation as described herein, including but not limited to a solution, suspension (incl. colloidal suspension), slurry, emulsion, homogenate, or the like. A preferred aqueous formulation is a solution, and therefore the term "solution" has been used in the exemplary sense throughout the detailed description of the preferred embodiments to refer to any of the aqueous compositions of the invention.

Specimen Collection for Clinical Diagnostic Laboratories

Collection is first step in diagnostic platforms or molecular protocols requiring the detection of potentially minute amounts of nucleic acids from pathogens including viruses. To facilitate the dynamic advancements in nucleic acid based detection strategies and their integration into the mainstream diagnostic laboratories there is a colossal need for reliable, robust, and standardized collection systems developed specifically with the intent of being utilized for downstream nucleic acid based detection such as the aforementioned platforms. The invention can alternatively be adapted for transport of nucleic acids from a doctor's office or operating room, or alternatively transported to a regional center, such as a hospital.

A clinical or veterinary specimen or a forensic or environmental sample collection system may include one or more collection tools and one or more reagents for efficiently: 1) obtaining a high yield of suitable specimen beyond what is currently available in the art; 2) inactivating potentially infectious biological pathogens so that they are no longer viable and can be handled; shipped, or transported with minimal fear of pathogen release or contamination; or 3) effectively stabilizing and preserving lysed `naked` RNA/DNA polymers from hydrolysis or nuclease degradation for prolonged periods at ambient temperatures until samples can be processed at a diagnostic laboratory, and preferably for achieving two or more, or all three, of these goals.

The collection/transport solutions of the present invention can provide a number of improvements and benefits over those presently available in the art. Exemplary benefits include, without limitation, one or more of the following:

Inactivation, killing, and/or lysis of microbes, viruses, or pathogens;

Destruction and/or inactivation of exogenous or endogenous nucleases, including, without limitation, RNase and/or DNase;

Compatibility with a variety of conventional nucleic acid extraction, purification, and amplification systems;

Preservation of RNA and/or DNA integrity within the sample;

Facilitation of transport and shipping at ambient temperatures, even over extended periods of time, or extreme temperature variations; and

Suitability for short--(several hours to several days), intermediate--(several days to several weeks), or long--(several weeks to several months) term storage of the isolated nucleic acids.

The disclosed compositions are particularly well suited for point-of-care, field studies, in-home health care or testing, triage/emergency and casualty assessment(s), mobile forensics, pathology, epidemiological sampling, crime scene investigation, paternity testing, pre- and post-pregnancy genetic screening, rape/incest testing and family counseling, confidential screening and testing for sexually transmitted diseases, including, without limitation, HIV, syphilis, Chlamydia, gonorrhoeae, or other venereal diseases and the like, and may be of particular value during the monitoring, etiology, and control of epidemic or pandemic diseases in both human and animal populations domestically and abroad. The compositions may be of particular relevance in collecting and analyzing Influenzavirus samples, including without limitation to predict and help manage shift and drift and to manage an imminent or ongoing pandemic.

In certain embodiments, the nucleic acid(s) isolated by the methods of the present invention may serve as a template in one or more subsequent molecular biological applications, assays, or techniques, including, without limitation, genetic fingerprinting; amplified fragment length polymorphism (AFLP) polymerase chain reaction (PCR); restriction fragment length polymorphism analysis (RFLP); allele-specific oligonucleotide analysis (ASOA); microsatellite analysis; Southern hybridization; Northern hybridization; variable number of tandem repeats (VNTR) PCR; dot-blot hybridization; quantitative real-time PCR; polymerase cycling assembly (PCA); nested PCR; quantitative PCR (Q-PCR); asymmetric PCR; DNA footprinting; single nucleotide polymorphism (SNP) genotyping; reverse transcription PCR(RT-PCR); multiplex PCR (m-PCR); multiplex ligation-dependent probe amplification (MLPA); ligation-mediated PCR (LmPCR); methylation specific PCR (MPCR); helicase-dependent amplification (HDA); overlap-extension PCR (OE-PCR); whole-genome amplification (WGA); plasmid isolation; allelic amplification; site-directed mutagenesis; high-throughput genetic screening; or the like, or any combination thereof.

The compositions of the present invention provide clinical/environmental collection solutions that efficiently achieve at least three, and preferably all four of the following: 1) kill or inactivate potentially-infectious pathogens, so that they are non-viable and can be safely handled, shipped or transported; 2) lyse cells to release RNAs and/or DNAs from the biological specimen contained in the collection system; 3) protect the released or `naked` polynucleotides in the sample from further hydrolysis or enzymatic degradation, modification, or inactivation; and 4) prolong the conventional time-frame for storage and transportation of the processed sample under a variety of ambient, sub- or supra-optimal temperature conditions to maintain the fidelity and integrity of the released polynucleotides until the biological material can be further processed or analyzed at a diagnostic facility or analytical laboratory.

In one exemplary embodiment, the methods and formulations maintain at least substantial stability of the nucleic acids in the sample for an extended period of time, e.g., for up to about 15 days, preferably up to about 30 days, or more preferably up to about 60 days or more, without refrigeration or freezing of the sample, and even when stored at room temperature, or ambient environmental conditions including those of temperate, sub-tropical or tropical climates and the like. In other embodiments, use of the disclosed compositions to prepare nucleic acids from a sample of biological origin is desirable to maintain at least substantial integrity and fidelity of the nucleic acids released from the sample for extended periods including, without limitation, at least about 5 to about 15 days, preferably at least about 10 to 20 days, more preferably at least about 15 to 25 days, or more preferably still, at least about 20 to 30 days or more, without a requirement for refrigerating or freezing the sample either at the time of sample collection or until the sample is further processed (or both) hours, days, weeks, or even months after originally being collected and placed into the disclosed storage/collection/transport/stabilization formulations.

Nucleic acids obtained from biological samples in the practice of the disclosed methods are advantageously compatible with a number of conventional molecular and diagnostic isolation, purification, detection, and/or analytic methodologies. The disclosed compositions facilitate recovery storage, and transport of populations of stabilized, substantially non-degraded, polynucleotides for use in a variety of subsequent methodologies, including, without limitation, nucleic acid isolation, purification, amplification, and molecular analytical and/or diagnostic testing, assay, analysis, or characterization, and the like.

Exemplary Commercial Kits of the Present Invention

The following outline provides exemplary commercial kits employing the PrimeStore.TM. compositions of the present invention (FIG. 5).

Peel-Pouch Collection System

Five-mL tube containing 1.5 mL PrimeStore.TM. Solution; Collection swab (e.g., FlockedSWABS.RTM. [Copan Diagnostics, Inc., Murrieta, Calif., USA]); and instructions for collection and/or processing of samples. (packed, e.g., in 50 pouches/unit) (See FIG. 4A, FIG. 4B, FIG. 4C, FIG. 4D, and FIG. 4E for a schematic demonstration of such systems).

PrimeStore.TM. Stock Solution (e.g., 25-mL bottles)

Once formulated, PrimeStore.TM. stock solution is stable at 4.degree. C. or below for periods of at least one year or more. Formulated PrimeStore.TM. Solution has also been shown to be stable at ambient temperature (e.g., about 20-30.degree. C.) for periods of three to six months or more.

Once a sample is contacted with a PrimeStore.TM. formulation as disclosed herein, it can reasonably expected to be stored indefinitely at temperatures of 0.degree. C. or below, at least one year or more under refrigeration (e.g., 4.degree. C. and at least 30 days or more at ambient temperature (e.g., about 20-30.degree. C.), without significant loss of nucleic acid composition, fidelity or integrity of the sample. For example, without limitation, the integrity of a population of polynucleotides obtained from the sample is at least substantially maintained, and preferably entirely maintained without detectable degradation, when the composition comprising the sample is stored at a temperature of from about -20.degree. C. to about 40.degree. C., for a period of from about 30 days to about 60 days.

The kit may also include one or more vials including the inventive compositions and one or more extraction devices to help liberate and separate the nucleic acids to obtain at least substantially pure RNA/DNA to be analyzed.

Environmental Sample and Storage Systems

0.1-, 0.2-, 0.5-, 1-, 2-, or 3-mL collection vials each containing 0.1 mL, 0.2 mL, 0.25 mL, 0.5 mL, 0.75 mL, or 1 mL PrimeStore.TM. solution; and instructions for collection and/or processing of samples (packed, e.g., 10 vials/unit). The collection vials may be sized larger as needed depending on the proposed collection method.

Definitions

The terms "about" and "approximately" as used herein, are interchangeable, and should generally be understood to refer to a range of numbers around a given number, as well as to all numbers in a recited range of numbers (e.g., "about 5 to 15" means "about 5 to about 15" unless otherwise stated). Moreover, all numerical ranges herein should be understood to include each whole integer within the range.

As used herein, the term "carrier" is intended to include any solvent(s), dispersion medium, coating(s), diluent(s), buffer(s), isotonic agent(s), solution(s), suspension(s), colloid(s), inert(s) or such like, or a combination thereof that is pharmaceutically acceptable for administration to the relevant animal or acceptable for a diagnostic purpose, as applicable. The use of one or more delivery vehicles for chemical compounds in general, and peptides and epitopes in particular, is well known to those of ordinary skill in the pharmaceutical arts. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the diagnostic, prophylactic, and therapeutic compositions is contemplated. One or more supplementary active ingredient(s) may also be incorporated into, or administered in association with, one or more of the disclosed immunogenic compositions.

As used herein, the term "nucleic acid" includes one or more types of: polydeoxyribonucleotides (containing 2-deoxy-D-ribose), polyribonucleotides (containing D-ribose), and any other type of polynucleotide that is an N-glycoside of a purine or pyrimidine base, or modified purine or pyrimidine bases (including abasic sites). The term "nucleic acid," as used herein, also includes polymers of ribonucleosides or deoxyribonucleosides that are covalently bonded, typically by phosphodiester linkages between subunits, but in some cases by phosphorothioates, methylphosphonates, and the like. "Nucleic acids" include single- and double-stranded DNA, as well as single- and double-stranded RNA. Exemplary nucleic acids include, without limitation, gDNA; hnRNA; mRNA; rRNA, tRNA, micro RNA (miRNA), small interfering RNA (siRNA), small nucleolar RNA (snoRNA), small nuclear RNA (snRNA), and small temporal RNA (stRNA), and the like, and any combination thereof.

As used herein, the terms "protein," "polypeptide," and "peptide" are used interchangeably, and include molecules that include at least one amide bond linking two or more amino acid residues together. Although used interchangeably, in general, a peptide is a relatively short (e.g., from 2 to about 100 amino acid residues in length) molecule, while a protein or a polypeptide is a relatively longer polymer (e.g., 100 or more residues in length). However, unless specifically defined by a chain length, the terms peptide, polypeptide, and protein are used interchangeably.

As used herein, "sample" includes anything containing or presumed to contain a substance of interest. It thus may be a composition of matter containing nucleic acid, protein, or another biomolecule of interest. The term "sample" can thus encompass a solution, cell, tissue, or population of one or more of the same that includes a population of nucleic acids (genomic DNA, cDNA, RNA, protein, other cellular molecules, etc.). The terms "nucleic acid source," "sample," and "specimen" are used interchangeably herein in a broad sense, and are intended to encompass a variety of biological sources that contain nucleic acids, protein, one or more other biomolecules of interest, or any combination thereof. Exemplary biological samples include, but are not limited to, whole blood, plasma, serum, sputum, urine, stool, white blood cells, red blood cells, buffy coat, swabs (including, without limitation, buccal swabs, throat swabs, vaginal swabs, urethral swabs, cervical swabs, rectal swabs, lesion swabs, abscess swabs, nasopharyngeal swabs, and the like), urine, stool, sputum, tears, mucus, saliva, semen, vaginal fluids, lymphatic fluid, amniotic fluid, spinal or cerebrospinal fluid, peritoneal effusions, pleural effusions, exudates, punctates, epithelial smears, biopsies, bone marrow samples, fluid from cysts or abscess contents, synovial fluid, vitreous or aqueous humor, eye washes or aspirates, pulmonary lavage or lung aspirates, and organs and tissues, including but not limited to, liver, spleen, kidney,

lung, intestine, brain, heart, muscle, pancreas, and the like, and any combination thereof. In some embodiments, the sample may be, or be from, an organism that acts as a vector, such as a mosquito, or tick, or other insect(s).

Tissue culture cells, including explanted material, primary cells, secondary cell lines, and the like, as well as lysates, homogenates, extracts, or materials obtained from any cells, are also within the meaning of the term "biological sample," as used herein. Microorganisms (including, without limitation, prokaryotes such as the archaeobacteria and eubacteria; cyanobacteria; fungi, yeasts, molds, actinomycetes; spirochetes, and mycoplasmas); viruses (including, without limitation the Orthohepadnaviruses [including, e.g., hepatitis A, B, and C viruses], human papillomavirus, Flaviviruses [including, e.g., Dengue virus], Lyssaviruses [including, e.g., rabies virus], Morbilliviruses [including, e.g., measles virus], Simplexviruses [including, e.g., herpes simplex virus], Polyomaviruses, Rubulaviruses [including, e.g., mumps virus], Rubiviruses [including, e.g., rubella virus], Varicellovirus [including, e.g., chickenpox virus], rotavirus, coronavirus, cytomegalovirus, adenovirus, adeno-associated virus, baculovirus, parvovirus, retrovirus, vaccinia, poxvirus, and the like), algae, protozoans, protists, plants, bryophytes, and the like, and any combination of any of the foregoing, that may be present on or in a biological sample are also within the scope of the invention, as are any materials obtained from clinical or forensic settings that contain one or more nucleic acids are also within the scope of the invention. The ordinary-skilled artisan will also appreciate that lysates, extracts, or materials obtained from any of the above exemplary biological samples are also within the scope of the invention.

As used herein, the term "buffer" includes one or more compositions, or aqueous solutions thereof, that resist fluctuation in the pH when an acid or an alkali is added to the solution or composition that includes the buffer. This resistance to pH change is due to the buffering properties of such solutions, and may be a function of one or more specific compounds included in the composition. Thus, solutions or other compositions exhibiting buffering activity are referred to as buffers or buffer solutions. Buffers generally do not have an unlimited ability to maintain the pH of a solution or composition; rather, they are typically able to maintain the pH within certain ranges, for example from a pH of about 5 to 7.

As used herein, the term "biological molecule" refers to any molecule found within a cell or produced by a living organism, including viruses. This may include, but is not limited to, nucleic acids, proteins, carbohydrates, and lipids. As used herein, a "cell" refers to the smallest structural unit of an organism that is capable of independent functioning and is comprised of cytoplasm and various organelles surrounded by a cell membrane. This may include, but is not limited to, cells that function independently such as bacteria and protists, or cells that live within a larger organism such as leukocytes and erythrocytes. As defined herein, a cell may not have a nucleus, such as a mature human red blood cell.

Samples in the practice of the invention can be used fresh, or can be used after being stored for a period of time, or for an extended period of time, including for example, cryopreserved samples and the like, and may include material of clinical, veterinary, environmental or forensic origin, may be isolated from food, beverages, feedstocks, potable water sources, wastewater streams, industrial waste or effluents, natural water sources, soil, airborne sources, pandemic or epidemic populations, epidemiological samples, research materials, pathology specimens, suspected bioterrorism agents, crime scene evidence, and the like.

As used herein, the term "patient" (also interchangeably referred to as "host" or "subject") refers to any host that can serve as a source of one or more of the biological samples or specimens as discussed herein. In certain aspects, the donor will be a vertebrate animal, which is intended to denote any animal species (and preferably, a mammalian species such as a human being). In certain embodiments, a "patient" refers to any

animal host, including but not limited to, human and non-human primates, avians, reptiles, amphibians, bovines, canines, caprines, cavines, corvines, epines, equines, felines, hircines, lapines, leporines, lupines, murines, ovines, porcines, racines, vulpines, and the like, including, without limitation, domesticated livestock, herding or migratory animals or birds, exotics or zoological specimens, as well as companion animals, pets, and any animal under the care of a veterinary practitioner. The invention may also be used to monitor disease outbreak, progression, and epidemiological statistics for a variety of global populations, including, without limitation, wasting disease in ungulates, tuberculosis, ebola, SARS, and avian influenzas. In certain embodiments, the samples will preferably be of mammalian origin, and more preferably of human origin.

The term "chaotrope" or "chaotropic agent" as used herein, includes substances that cause disorder in a protein or nucleic acid by, for example, but not limited to, altering the secondary, tertiary, or quaternary structure of a protein or a nucleic acid while leaving the primary structure intact.

The term "e.g.," as used herein, is used merely by way of example, without limitation intended, and should not be construed as referring only those items explicitly enumerated in the specification.

The term "substantially free" or "essentially free," as used herein, typically means that a composition contains less than about 10 weight percent, preferably less than about 5 weight percent, and more preferably less than about 1 weight percent of a compound. In a preferred embodiment, these terms refer to less than about 0.5 weight percent, more preferably less than about 0.1 weight percent or even less than about 0.01 weight percent. The terms encompass a composition being entirely free of a compound or other stated property, as well. With respect to degradation or deterioration, the term "substantial" may also refer to the above-noted weight percentages, such that preventing substantial degradation would refer to less than about 15 weight percent, less than about 10 weight percent, preferably less than about 5 weight percent, etc., being lost to degradation. In other embodiments, these terms refer to mere percentages rather than weight percentages, such as with respect to the term "substantially non-pathogenic" where the term "substantially" refers to leaving less than about 10 percent, less than about 5 percent, etc., of the pathogenic activity.

In accordance with long standing patent law convention, the words "a" and "an" when used in this application, including the claims, denotes "one or more."

EXAMPLES

The following examples are included to demonstrate illustrative embodiments of the invention. It should be appreciated by those of ordinary skill in the art that the techniques disclosed in the examples that follow represent techniques discovered to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of ordinary skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1

Formulation of Exemplary Storage Solutions

The present example provides a general formulation of the PrimeStore.TM. (PanFlu) compositions of the present invention. Exemplary formulations are also detailed in Examples 2-5.

Materials

Guanidine thiocyanate, sodium citrate, Antifoam A.RTM. Concentrate, and N-lauroylsarcosine, sodium salt, were all purchased from Sigma Chemical Co. (St. Louis, Mo., USA). Tris(2-carboxyethyl) phosphine hydrochloride (TCEP) was obtained from Soltec Ventures Inc. (Beverly, Mass., USA). 2-amino-2-hydroxymethyl-propane-1,3-diol (TRIS) was obtained from Applied Biosystems/Ambion (Austin, Tex., USA). 2-[2-(Bis(carboxymethyl)amino)ethyl-(carboxymethyl)amino]acetic acid (EDTA) GIBCO.RTM. Ultra Pure was obtained from Invitrogen Corp. (Carlsbad, Calif., USA). All other reagents are available commercially from Sigma-Aldrich or USB Corporation.

TABLE-US-00001 TABLE 1 FORMULATION RANGES OF EXEMPLARY COMPONENTS FOR THE PREPARATION OF PRIMESTORE .TM. COMPOSITIONS

Reagent	Component	Final Concentration
1. A chaotrope, e.g.: Guanidine thiocyanate	about 0.5M to about 6M	or Guanidine hydrochloride about 0.5M to about 6M
2. An anionic detergent, e.g.: N-lauroyl sarcosine	about 0.15% to about 1% (wt./vol.)	(inter alia Na salt) or Sodium dodecyl sulfate, Same Lithium dodecyl sulfate, Same Sodium glycocholate, Same Sodium deoxycholate, Same Sodium taurodeoxycholate, or Same Sodium cholate about 0.1% to about 1% (wt./vol.)
3. A reducing agent, e.g.: TCEP	about 0.5 mM to about 30 mM	or .beta.-ME, DTT, formamide, about 0.05M to about 0.3M
4. A chelator, e.g.: Sodium citrate	about 0.5 mM to about 50 mM	or EDTA, EGTA, HEDTA, about 0.01 mM to about 1 mM
5. A buffer (e.g., TRIS, HEPES, MES, Bis-Tris, etc.)	about 1 mM to about 1M	MOPS, etc.)
6. An acid q.s. to adjust to a pH of about 6 to 7, (e.g., HCl or citric acid)	preferably 6.4 to 6.8	
7. Nuclease-free water q.s. to desired final volume	Optionally one or more of:	
8. A surfactant/defoaming agent, about 0.0001% to about 0.3% (wt./vol.)	e.g.: Antifoam A .RTM. or Tween .RTM.	
9. An alkanol (e.g., methanol, about 1% to about 25% (vol./vol.) ethanol, propanol, etc.)		
10. RNA or DNA	about 1 pg to about 1 .mu.g/mL	

Example 2

Formulation of an Exemplary Storage Solution

The present example describes a first exemplary formulation of the compositions of the invention. This formulation has also been alternatively referred to by the inventors as "PrimeStore.TM. Solution" or "PSS" version 1.

TABLE-US-00002 TABLE 2 PREPARATION OF PRIMESTORE .TM. COMPOSITION (VER. 1)

Reagent	Final Concentration
Guanidine thiocyanate	4M
Sodium citrate	30 mM
Sodium dodecyl sulfate	0.25% (wt./vol.)
N-lauroyl sarcosine, sodium salt	0.25% (wt./vol.)
2-mercaptoethanol (.beta.-ME)	0.1M
Antifoam A	0.1% (wt./vol.)
Citric acid q.s. to adjust pH to 6.5	
Nuclease-free water	11.82 mL

Example 3

Preparation of a Second Exemplary Storage Solution

The present example describes the preparation of another exemplary storage solution according to the present invention. This formulation has also been alternatively referred to by the inventors as PrimeStore.TM. version 2.

TABLE-US-00003 TABLE 3 PREPARATION OF PRIMESTORE .TM. COMPOSITION (VER. 2) Reagent Quantity Final Concentration Guanidine thiocyanate 35.488 gm 3M TCEP 0.02867 gm 1 mM Sodium citrate 0.2931 gm 10 mM N-lauroyl sarcosine, 0.5 gm 0.5% sodium salt (NLS) Antifoam A (10% solution) 200 .mu.L 0.002% TRIS (1M) 10 mL 100 mM EDTA (0.5M) 20 .mu.L 0.1 mM Hydrochloric acid (HCl) q.s. to adjust pH to 6.7 -- Nuclease-free water q.s. to 100 mL --

Example 4

Preparation of a Third Exemplary Storage Solution

The present example describes the preparation of another exemplary storage solution according to the present invention. This formulation has also been alternatively referred to by the inventors as PrimeStore.TM. version 2.2.

TABLE-US-00004 TABLE 4 PREPARATION OF PRIMESTORE .TM. COMPOSITION (VER. 2.2) Reagent Quantity Final Concentration Guanidine thiocyanate 35.488 gm 3M TCEP 0.02867 gm 1 mM Sodium citrate 0.2931 gm 10 mM N-lauroyl sarcosine, sodium salt (NLS) 0.5 gm 0.5% Antifoam A (10% solution) 200 .mu.L 0.002% TRIS (1M) 10 mL 100 mM EDTA (0.5M) 20 .mu.L 0.1 mM Ethanol, molecular grade (96-100%) 23 mL 23% (vol./vol.) Hydrochloric acid (HCl) q.s. to adjust pH to 6.7 -- Nuclease-free water q.s. to 100 mL -- Exemplary Protocol for Preparation of PrimeStore Solution (ver. 2.2) 1. Add 40 mL of nuclease-free water to a clean beaker with a stir bar. 2. Place beaker on a hot plate/stirrer and adjust temperature to 60-65.degree. C. Set stirring speed to medium. 3. Add 35.488 gm of guanidine thiocyanate slowly to the water allowing it to dissolve as added. 4. Add 0.0287 gm of TCEP to beaker and increase stirrer speed to help dissolve crystals. 5. Add 0.2931 gm of sodium citrate to the beaker. 6. Add 0.5 gm of NLS to the solution. Increase stirrer speed to create a vortex in the beaker. This will pull the NLS into the solution and help dissolve the reagent. 7. Vortex a prepared 10% Antifoam A solution (1 mL Antifoam A Concentrate + 9 mL nuclease-free water). Pipette 200 .mu.L of the 10% Antifoam A into the solution. 8. Pipette 10 mL of 1M TRIS into the solution. 9. Pipette 20 .mu.L of 0.5M EDTA into the solution. 10. Increase the temperature to bring the solution to 75-80.degree. C. and stir for 3-5 minutes. 11. Remove beaker from heat and allow solution to cool to room temperature (.apprxeq.22-25.degree. C.). 12. Add 23 mL of ethanol to the solution and mix thoroughly. 13. Adjust pH to 6.7 with HCl. 14. Pour solution into a clean 100 mL graduated cylinder. 15. Add nuclease-free water to bring total volume to 100 mL. 16. Transfer solution to a labeled sterile container. Store at room temperature (.apprxeq.22-25.degree. C.). *Note: Preferably, make sure each reagent is completely dissolved before adding the next.

Example 5

Comparison of PrimeStore.TM. Solutions to Conventional Formulations

A sample of homogenized nasal tissue from a cotton rat (*Sigmodon hispidus*) challenged with influenza A (H3N2) or a human clinical influenza A (H1N1) sample collected as a human clinical nasal wash during the 2006-07 season were placed in PrimeStore.TM. Solution (Ver. 1) and tested compared to the respective lysis formulation and protocol, and extraction procedure, from three commercially available kits: RNAqueous.RTM.-Micro (Ambion, Austin, Tex., USA), QIAamp Viral RNA Mini Kit (Catalogue #52904, Qiagen, Valencia, Calif., USA), and MagMax AI/ND Viral RNA Isolation Kit (Catalogue #AM1929, Ambion). Extraction efficiency was evaluated using the ABI 7500 sequence detection system with the

comparative threshold cycle (C.sub.T) method (See FIG. 2). In FIG. 2, "delta Rn" represents the fluorescent reporter signal minus a baseline amount. As shown in FIG. 1 and FIG. 2, the relative C.sub.T scores and viral copies detected were optimal when the fixing formulation was used in place of the respective conventional lysis buffer for each commercial kit. In these two sample types, the compositions of the invention worked better than the two conventional Kits for extraction purposes. The PrimeStore.TM. Solution (ver. 1) composition was also shown to be readily compatible with commercially available nucleic acid extraction kits. FIG. 1 illustrates RNA extraction results where the version 1 of PrimeStore.TM. Solution was used in conjunction with three commercially available kits: Qiagen Viral Mini, Ambion RNAqueous Mini, and Ambion A1/NCD MagMax. As illustrated by FIG. 1, when the lysis buffer of the extraction kit was replaced with the fixing formulation (denoted on the figure as "One-Step+), superior nucleic acid extraction was achieved when compared to extraction using kits according to standard protocol (denoted on the figure as "One-Step-". Extraction efficiency was measured by real time (r) reverse transcription (RT) polymerase chain reaction (PCR) [rRT-PCR].

FIG. 3 shows preservation of naked RNA in PrimeStore.TM. Solution compared to preservation in a prior solution, with water used as a control. As illustrated in FIG. 3, detection (by fluorescence) occurred at the earliest amplification cycle for RNA stored in PrimeStore.TM. Solution (ver. 1) at all time-points assayed.

Example 6

PrimeStore.TM. Solution for the Collection of Nasal Wash Specimens

A prospective clinical detection study was conducted using nasal wash specimens from: 1) symptomatic pediatric patients and 2) asymptomatic or symptomatic family members. Detection of influenza virus compared nasal wash specimens collected in PrimeStore.TM. Solution and Viral Transport Medium (VTM) by real-time RT-PCR (rRT-PCR) and traditional culture, respectively. Genetic characterization of influenza genes encoding hemagglutinin (HA), neuraminidase (NA), and matrix surface (MA) proteins were performed using select nasal wash specimens preserved in PrimeStore.TM. Solution to evaluate vaccine effectiveness and drug sensitivity within viral strains.

Influenza is a highly evolving, RNA-based respiratory virus responsible for more than 200,000 hospitalizations and about 36,000 fatalities each year in the United States. Widespread emergence of influenza drift variants among contemporary circulating human viruses prompted a change in all three vaccine components for the upcoming 2008/09 season. Increased morbidity and mortality during the 2007/08 season included 72 influenza-associated pediatric deaths and continued drug resistance (oseltamivir [TamiFlu.RTM., Roche Laboratories, Inc., Nutley, N.J., USA] and adamantadine) within circulating strains.

Materials and Methods

A total of 100 pediatric (index) patients who met the clinical case criteria for influenza infection and 126 family contacts were enrolled in the study. Nasal washings were placed into PrimeStore.TM. Solution and Universal Viral Transport Medium (Becton-Dickinson, Franklin Lakes, N.J., USA) and analyzed by rRT-PCR or culture analysis, respectively. rRT-PCR was performed using influenza type (A or B) and subtype (H3, H1, H5) specific primers/probes according to Daum et al. (2007). Further genetic characterization of selected clinical samples preserved in PrimeStore.TM. Solution was performed using standard RT-PCR and direct nucleotide sequencing of the hemagglutinin HA, NA, and MA viral proteins.

Results

Of the total samples evaluated (N=226; 100 index, 126 family contacts), 66 (29%) tested positive for influenza virus (45 H3N2, 2 H1N1 and 19 B) by rRT-PCR. rRT-PCR from nasal washings preserved in PrimeStore.TM. Solution detected influenza virus from 11 patients (9 Flu A and 2 Flu B) that were not detected by culture (Table 5 and Table 6). Of these 11 specimens, five were from patients enrolled as family contacts.

Phylogenetic analysis of influenza A and B HA genes exhibited drifting compared to the 2007/08 vaccine strains and revealed a higher genetic homology to the 2008/09 Brisbane vaccine strains. Some genetic differences in viruses were noted among family members, particularly among influenza A (H3N2) strains. MA analysis revealed adamantane resistance in all influenza A H3N2 strains, but sensitivity in both H1N1 viruses. All influenza B strains (n=18) were sensitive to the neuraminidase inhibitor drugs zanamivir (Relenza.RTM. GlaxoSmithKline, Research Triangle Park, N.C., USA) and oseltamivir (Tamiflu.RTM. Roche) based on the presence of an aspartic acid (D) at amino acid 197 (influenza B numbering) in the NA gene.

Real-Time RT-PCR vs. Culture

rRT-PCR is superior to traditional culture for the detection of influenza virus from original nasal wash specimens preserved in PrimeStore.TM. solutions: influenza was detected within 2 hours (c.f. 2 to 7 days for conventional culture methods); and the analyses were more sensitive (11 specimens; 9 Flu A and 2 Flu B detected below culture limits). Moreover, the use of molecular diagnostic methods in lieu of conventional organism culture did not propagate potentially infectious viruses, and simultaneously provided the type and subtype of the influenza virus.

Genetic Analysis

Vaccine Relatedness

H3N2 Strains: Analysis of the HA1 gene of the influenza A (H3N2) hemagglutinin (HA) revealed genetic drift including five amino acid differences in all Texas strains compared to the 2007-08 A/Wisconsin/67/2005 vaccine strain. One HA1 mutation noted in all the Texas strains (D122N) is a potential glycosylation site. All A/Texas (H3N2) strains exhibited a greater HA homology (99.0-99.7%) to the newly selected 2008-09 A/Brisbane/10/2007 strain.

H1N1 Strains: The hemagglutinin HA1 gene of the 2 influenza A (H1N1) exhibited 7 amino acid changes compared to A/Solomon Island/3/2006 vaccine strain. Four substitutions (R90K, T145V, K210T and E290K) were within known H1 antibody combining sites. Both Texas H1N1 strains exhibited greater HA homology (98.8% and 99.4%) to the newly selected 2008-09 A/Brisbane/59/2007 vaccine strain.

Influenza B strains: Analysis of the HA1 hemagglutinin and neuraminidase genes revealed all Texas strains were of the B/Yamagata lineage and genetically more homologous to the 2008-09 B/Brisbane/5/2007 vaccine strain than the 2007/08 B/Malaysia/2506/2004 vaccine strain.

Family Mutation

Amino acid changes were noted in the NA, HA1, M1 and M2e among family members. The HA1. Hemagglutinin showed the highest mutation of the influenza genes analyzed, with one family exhibiting five amino acid changes.

Analysis of the highly conserved 24 amino acid M2e viral surface proton pump showed some variation within families. One index patient strain contained 3 unique amino acid M2e substitutions that were `wild-type` within family member strains.

Antiviral Susceptibility

Adamantane: Matrix gene (MA) genetic analysis, specifically a serine-to-asparagine substitution at position 31 (S31N), revealed adamantane resistance in all influenza A (H3N2) strains but sensitivity in both influenza A (H1N1) viruses.

Neuraminidase Inhibitors: All Texas influenza A (H3N2) isolates were shown to be sensitive to oseltamivir through genetic analysis of E119V, R292K, and N294S substitutions in the NA gene. Genetic analysis of the influenza B NA gene revealed that all Texas strains contained an aspartic acid (D) residue at position 197, and are thus likely sensitive to oseltamivir.

The protocols and tests herein can be adapted for other microbes like tuberculosis, malaria, staphylococcus, and the like and other pathogens where there is a need to know antimicrobial susceptibility quickly.

Example 7

Influenza Sample Collection Using PrimeStore.TM. Solution

The compositions of the present invention provide a single sample collection, transport, and storage reagent that facilitate: 1) procuring high quality nucleic acids from clinical or environmental specimens, 2) inactivation of potentially infectious biological pathogens for safe handling and transport of specimens, and 3) stabilization and preservation of released `naked` RNA/DNA preventing hydrolysis/nuclease degradation for prolonged periods at ambient temperatures. The results of one such study are presented in the following example:

TABLE-US-00005 TABLE 5 Influenza Subtype Detection: rRT-PCR vs. Culture Total Flu A Samples
 Influenza A Influenza B Total Flu B Samples (N = 47) rRT-PCR (N = 47) Culture (N = 40) rRT-PCR (N = 19)
 Culture (N = 17) (N = 19) Index Patients (28) 28/28 (100%) 23/28 (82%) 16/16 (100%) 14/16 (88%) Index
 Patients (16) Family Contacts (19) 19/19 (100%) 15/19 (79%) 3/3 (100%) 3/3 (100%) Family Contacts (3)

TABLE-US-00006 TABLE 6 Positive Influenza Detection: rRT-PCR vs. Culture Total Samples Flu Positive
 rRT-PCR (N = 226) (N = 66) (N = 66) Culture (N = 66) Index Patients (100) 44/100 39/39 (100%) 37/39
 (94%) Family Contacts (126) 22/126 27/27 (100%) 18/27 (67%)

This example illustrates the effectiveness of the PrimeStore.TM. Solution (ver. 2.2) in killing pathogenic microbe(s).

Methods

Real-time RT-PCR was used to assay influenza A (H5N1) virus nucleic acid preserved in PrimeStore.TM. Solution. A time-course study at room temperature was carried out to evaluate the integrity of clinical specimens, cloacal samples, and cloned template avian influenza A virus (H5) RNA stored and extracted from PrimeStore.TM. Solution, Viral Transport Media, RNA Storage Solution, or nuclease-free water. PrimeStore.TM. Solution extraction efficiency was compared to three commercially available nucleic acid extraction kits. Furthermore, the ability of RNA contained in PrimeStore.TM. Solution to resist nuclease degradation was evaluated.

Results

PrimeStore.TM. Solution (version 2, but lacking ethanol) inactivated microbial agents while preserving released RNA/DNA from clinical material, i.e., nasal washes, throat swabs, or environmental samples. Clinical specimens or environmental samples placed in this solution were stabilized at room temperature for up to 30 days while degradation of nucleic acids occurred in other transport media. PrimeStore.TM. Solution is compatible with commercially available RNA isolation kits and produced an increased nucleic acid yield.

Example 8

Killing of MRSA (ATCC33592) in PrimeStore.TM. Solution

This example illustrates the effectiveness of the PrimeStore.TM. Solution (ver. 2.2) in killing a potential bacterial contaminant. Methicillin-resistant *Staphylococcus aureus* (MRSA) strain ATCC33592 was diluted 10-fold and 1000-fold into PrimeStore.TM. Solution (Ver. 2.2) and quantitated (see FIG. 6).

Experimental Protocol

Day Procedure

0 Transfer MRSA (ATCC33592) from a Culti-loop.RTM. (Remel) to 1.5 mL of TSB in a 15-ml conical test tube. Incubate at 37.degree. C. for approximately 15 min. Gently vortex suspension and transfer 100 .mu.L to a blood-agar plate. Incubate the plate overnight at 37.degree. C. 1 Observe heavy and uniform colony growth after 12 hr incubation. Transfer .about.10% of colonies to 300 mL of tryptic soy broth (TSB) in a sterile, 1-liter flask. Place flask on shaker at 37.degree. C. and 200 rpm.

After approximately 4-6 hrs' incubation, transfer .about.50 mL of bacterial suspension to new 1-liter flask containing 300 mL fresh TSB.

After approximately 4-6 hrs' incubation, transfer .about.100 .mu.L of culture into 900 .mu.L of TSB (1:10 control dilution). From this suspension 10 were transferred to 990 .mu.L of TSB (1:1000 control dilution).

Transfer 100 .mu.L, of culture into 900 .mu.L of PrimeStore.TM. Solution (1:10 PrimeStore.TM. dilution). From this suspension 10 .mu.L were transferred to 990 .mu.L of TSB (1:1000 PrimeStore.TM. dilution).

Immediately after transfer into TSB or PrimeStore.TM. Solution (ver. 2.2), the suspensions were gently vortexed and 100 .mu.L were plated from both dilutions of control and PrimeStore.TM. suspensions onto blood agar plates. The time-zero time point was actually about two minutes following addition of the bacteria to the TSB or PrimeStore.TM. Solution.

The suspensions in TSB and PrimeStore.TM. Solution (ver. 2.2) were kept at room temperature.

An additional 100 .mu.L was plated onto blood agar plates at 5, 15, 30, 60, 120 and 240 minutes after the preparation of the suspensions in TSB and PrimeStore.TM. Solution.

Bacterial suspensions on the plates were allowed to dry, the plated inverted and the plates incubated overnight at 37.degree. C.

A titration of the shaker culture was also performed by mixing 100 .mu.L of the suspension from the shaker culture with 900 .mu.L of TSB (10.sup.-1 dilution). Serial 10-fold dilutions were prepared through 10.sup.-9. 100 .mu.L samples were plated onto blood agar from all dilutions except 10.sup.-1.

Plates were observed for bacterial colonies. All plates were stored at 4.degree. C. for later observation, if necessary.

Results

The results are presented in Table 7 and Table 8. Briefly, the bacterial suspension contained approximately 4.7.times.10.sup.9 cfu/ml. Thus, the 1:10 dilution contained approximately 4.7.times.10.sup.8 cfu/ml and the 1:1000 dilution contained 4.7.times.10.sup.6 cfu/ml. At all time points, the bacteria suspended in TSB were too numerous to count. At all time points the bacteria suspended in PrimeStore.TM. Solution and plated onto blood agar plates had no detectable colonies.

TABLE-US-00007 TABLE 7 KILLING OF MRSA (ATCC 33592) BY PRIMESTORE .TM. SOLUTION (VER. 2.2) Incubation Time In TSB In Primestore (Minutes) 1:10 1:1000 1:10 1:1000 0 TNTC TNTC 0 0 5 TNTC TNTC 0 0 15 TNTC TNTC 0 0 30 TNTC TNTC 0 0 60 TNTC TNTC 0 0 120 TNTC TNTC 0 0 240 TNTC TNTC 0 0 TNTC = too numerous to count.

TABLE-US-00008 TABLE 8 TITRATION OF MRSA ATCC33592 FROM SUSPENSION CULTURE Dilution CFU/plate CFU/ml 1.E+01 TNTC 1.E+02 TNTC 1.E+03 TNTC 1.E+04 TNTC 1.E+05 TNTC 1.E+06 TNTC 1.E+07 35 3.5 .times. {circumflex over (10)}9 1.E+08 6 6 .times. {circumflex over (10)}9 1.E+09 0 NOTE: CFU/ml calculations are corrected to include the plating volume of 0.1 mls Final Conc: 4.7 .times. 10{circumflex over ()}9/ml TNTC = too numerous to count. CFU = colony forming units.

An additional study was performed to determine the time of exposure necessary for killing MRSA ATCC33592 when diluted 10-fold into PrimeStore.TM. Solution (Ver. 2.2), and to determine the effect of dilution of the bacteria after exposure to PrimeStore.TM. Solution, but before plating.

Experimental Protocol

Day Procedure

0 Transfer MRSA (ATCC33592) from TNTC plate from the study described above into 4 mL of TSB. These plates had been stored at 4.degree. C. for approximately 48 hrs. Bacteria were vortexed gently and placed at room temperature for approximately 10 min. before use. 0.1 mL of bacterial suspension was transferred to 0.9 mL PrimeStore.TM. Solution and vortexed gently. After approximately 60 sec, the bacteria in

PrimeStore.TM. were again vortexed gently and 0.1 mL of bacterial suspension was transferred into 0.3 mL of TSB (1:4 dilution). 100 .mu.L, of bacteria in PrimeStore.TM. Solution (designated "neat") and from the 1:4 dilution into TSB were plated onto blood agar plates (5% sheep RBCs in TSA). This process was repeated at 5 and 15 min., and then again with dilutions made into TSB instead of PrimeStore.TM. Solution.

The liquid bacterial suspensions on the blood agar plates were allowed to dry at room temperature and then incubated overnight at 37.degree. C.

1 After approximately 16 hrs. incubation, the plates were removed from the incubator and colonies counted.

Results

The bacterial suspension contained an unknown number of colony forming units (cfu) per mL. At all time points the bacteria suspended in tryptic soy broth (TSB) were too numerous to count (TNTC). At all timepoints the bacteria suspended in PrimeStore.TM. compositions and plated onto blood agar plates produced no colonies (Table 9).

TABLE-US-00009 TABLE 9 KILLING OF MRSA (ATCC33592) BY PRIMESTORE .TM. SOLUTION
Incubation Time In TSB In TSB (minutes) neat 1:4 neat 1:4 1 TNTC TNTC 0 0 5 TNTC TNTC 0 0 15 TNTC
TNTC 0 0 TNTC = too numerous to count.

Example 9

Additional Studies Evaluating PrimeStore.TM. Solutions

The data in FIG. 7B illustrate the ability of PSS to inactivate microbes. Shown is a study in which chicken cloacal specimens were collected in PrimeStore.TM. Solution (Ver. 1). PrimeStore.TM. Solution inactivated the microbial agents in .ltoreq.1 hr. Four original chicken cloacal samples were immersed in PrimeStore.TM. Solution or water and subsequently plated on blood agar plates. These results showed that the disclosed composition could quickly kill or inactive microorganisms in the sample.

The data in FIG. 7C illustrate the ability of PSS to inhibit RNA base hydrolysis for 30 days at room temperature. RNA was incubated at room temperature (22-26.degree. C.) in PrimeStore.TM. (gel lane 1 and 3) and water (gel lane 2 and 4), and subsequently RT-PCR amplified (1500 base pairs) at Day 0 and Day 30. PrimeStore.TM. preserved collected RNA, and prevented RNA/DNA degradation at room temperature up to 30 days (see also, e.g., Table 11).

Flu Inhibition Assay

The reagents for this assay include

Trypsin Medium:

45 mL Sterile N/C EMEM

3 mL stock 7.5% Na Bicarbonate (2%)

1.5 mL SPG (1%)

75 .mu.L Trypsin (0.05%)

1.5 mL Fungizone (1%)

150 .mu.L Gentamicin

Filter medium.

Crystal Violet: 150 ml glutaric dialdehyde

2 gm crystal violet

2850 mL deionized water

Protocols Preparation of Serum Samples for Assay

Thaw and vortex serum samples. For each sample, label the lid of a corresponding Spin-X tube. Combine 450 .mu.L non-complete EMEM with 50 .mu.L serum into a Spin-X tube. Warm tubes containing the sera and EMEM in a 37.degree. C. water bath for 30 min. Centrifuge tubes at 8000 RPM for 2 min. at room temperature. Label and place samples into a -20.degree. C. freezer until assayed.

Dilution Plates

Load 160 .mu.L of each neat compound or serum sample into wells A1 through A12. Load the remaining wells with 120 .mu.L, trypsin medium. Using a multi-channel pipette, draw 40 .mu.L, of neat sample from row A and dilute into the corresponding wells in row B. Repeat dilution for each row, mixing well after each transfer. At row H, after mixing the transfer from row G, draw up 40 .mu.L, from each well and discard. Obtain virus stock (10⁶ TCID₅₀) from -80.degree. C. freezer and thaw. Dilute virus stock in trypsin media to a 10³ dilution. After serial dilutions are completed, transfer 120 .mu.L of influenza virus (10⁴ TCID₅₀ per ml) to all wells in the dilution plate. This results in a total of 240 .mu.L in all wells. Incubate dilution plate(s) at room temperature for 1 hour.

MDCK Cell Plates

Sterilize and place glass reservoir, comb dispenser and tubing inside the fume hood. Inside the hood, connect the tubing to the reservoir and fill nozzle of the comb. Connect the aspirator tube to the vacuum nozzle on the comb. Place the reservoir on an elevated surface and turn on the aspirator. Put PBS into the reservoir (1 L or more may be needed depending on the number of plates). Wash the cell plates 3.times. with the PBS comb (aspirate the medium, then press the button for roughly 1 second to wash the wells, repeat twice). Using a multi-channel pipette, transfer 50 .mu.L from each well in column 1 of the dilution plate to columns 1 through 4 of the cell plate. Transfer 50 .mu.L from each well in column 2 of the dilution plate to columns 5 through 8 of the cell plate. Transfer 50 .mu.L from each well in column 3 to columns 9 through 12 of the cell plate. Repeat transfer to additional cell plates for remaining samples. Incubate cell plates for 1 hour at 37.degree. C. After incubation period, add 50 .mu.L trypsin medium to all wells of the cell plates. Return plates to incubation chamber, and incubate for 4 days post-infection.

Staining

Add 100 .mu.L of Crystal Violet to all wells. Let sit for 1 hour. Rinse plates in cold running water and air dry.

TABLE-US-00010 TABLE 10 5 mM 10 mM 25 mM 35 mM 50 mM TITRATION OF TCEP USING WHOLE INFLUENZA A VIRUS 30.353 24.58 24.52 24.14 25.9582 30.2261 22.74 24.26 22.74 26.0337 30.28955 23.66 24.39 23.44 25.99595 0.089732 1.301076 0.183848 0.989949 0.053387 Titration of TCEP Using H5 Avian ssRNA 27.2 25.25 25.63 27.3 28.3039 26.73 24.89 25.36 27.62 26.6854 26.965 25.07 25.495 27.46 27.49465 0.33234 0.254558 0.190919 0.226274 1.144452

Time-Course Study of the Long-Term Stability of PrimeStore Compositions

The following data demonstrate the effectiveness of various PrimeStore compositions in preserving nucleic acid integrity over a thirty-day period with samples stored at room temperature. PrimeStore compositions have been compared to water alone, ethanol alone, commercial buffers such as VTM and AVL.

TABLE-US-00011 TABLE 11 30-DAY TIME-COURSE COMPARISON STUDY OF VARIOUS COMPOSITIONS DAY 1 PS-V1 PS-V1 PS-V2.2 VTM Water EtOH AVL (Year old) (new lot) PS-V2 (w/EtOH) PS-V1 PS-V1 PS-V2.2 VTM Water EtOH AVL (Year old) (new lot) PS-V2 (w/EtOH) 27.0225 26.1403 18.4463 24.2698 24.2607 23.9524 23.4426 20.2102 24.42 25.6044 18.3206 24.4789 24.3716 23.9615 23.7387 20.063 25.72125 25.87235 18.4463 24.37435 24.31615 23.95695 23.59065 20.1366 AVG 1.840245 0.378939 0.088883 0.147856 0.0784181 0.006435 0.209374 0.1040861- 2 STDEV DAY 6 PS-V1 PS-V1 PS-V2.2 VTM Water EtOH AVL (Year old) (new lot) PS-V2 (w/EtOH) 29.1988 29.3053 27.4058 37.9226 27.2379 27.165 24.53 22.4887 28.6799 28.7916 27.0781 40 26.4857 26.7658 24.4418 22.4676 28.93935 29.04845 27.24195 38.9613 26.8618 26.9654 24.4859 22.47815 AVG 0.366918 0.363241 0.231719 1.468944 0.5318857 0.282277 0.062367 0.01491995 STDEV DAY 12 PS-V1 PS-V1 PS-V2.2 VTM Water EtOH AVL (Year old) (new lot) PS-V2 (w/EtOH) 27.997 28.151 26.9011 40 30.8352 31.0478 25.8926 22.2074 28.0062 28.2211 26.2139 38.0439 30.4502 30.1935 25.3037 22.0025 28.0016 28.18605 26.5575 39.02195 30.6427 30.62065 25.59815 22.10495 AVG 0.006505 0.049568 0.485924 1.383172 0.2722361 0.604081 0.416415 0.14488618 STDEV DAY 20 PS-V1 PS-V1 PS-V2.2 VTM Water EtOH AVL (Year old) (new lot) PS-V2 (w/EtOH) 27.9851 28.7713 27.1105 40 30.1844 27.193 25.7407 20.8364 28.4067 27.7929 27.0105 40 30.2465 27.2274 25.6213 20.2843 28.1959 28.2821 27.0605 40 30.21545 27.2102 25.681 20.56035 AVG 0.298116 0.691833 0.070711 0 0.0439113 0.024324 0.084429 0.39039365 STDEV DAY 30 PS-V1 PS-V1 PS-V2.2 VTM Water EtOH AVL (Year old) (new lot) PS-V2 (w/EtOH) 29.23 31.9168 33.012 40 29.1993 30.2386 23.0589 20.9348 29.9067 31.3252 32.3001 40 28.827 29.6081 22.9662 20.4973 29.56835 31.621 32.65605 40 29.01315 29.92335 23.01255 20.71605 AVG 0.478499 0.418324 0.503389 0 0.2632559 0.445831 0.065549 0.30935922 STDEV- PS-V1 (year old) = One-year old PrimeStore Formulation (Ver. 1). PS-V1 (new lot) = Fresh PrimeStore Formulation (Ver. 2). PS-V2 = Fresh PrimeStore Formulation (Ver 2) (without ethanol). PS-V2.2 (w/EtOH) = Fresh PrimeStore Formulation (Ver. 2.2) (i.e. with ethanol).

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of exemplary embodiments, it will be apparent to those of ordinary skill in the art that variations may be applied to the composition, methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More

specifically, it will be apparent that certain agents that are both chemically- and physiologically-related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those of ordinary skill in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims. Accordingly, the exclusive rights sought to be patented are as described in the claims below.

* * * * *

